

Original Research Article

<https://doi.org/10.20546/ijcmas.2026.1506.017>

Haematological Alterations Associated with Naturally Occurring Nephropathogenic Infectious Bronchitis in Broiler Chickens

Sakshi Patel¹*, Biswajit Dutta¹, Pankaj Deka², S. A. Begum¹, Rupam Dutta³,
Abhijit Deka¹, Suchand Doloi² and Ajay Mendake⁴

¹Department of Veterinary Pathology, College of Veterinary Science, AVFU, Khanapara, Guwahati, India-781022

²Department of Veterinary Microbiology, College of Veterinary Science, AVFU, Khanapara, Guwahati, India-781022

³Department of Animal Biotechnology, College of Veterinary Science, AVFU, Khanapara, Guwahati, India-781022

⁴Department of Livestock Production and Management, Nagpur Veterinary College, MAFSU, Nagpur, India-440006

*Corresponding author

Keywords

Nephropathogenic
Infectious
Bronchitis,
Haematological
alterations,
Renal lesions,
Visceral uricosis

Article Info

Received:
15 April 2026
Accepted:
31 May 2026
Available Online:
10 June 2026

ABSTRACT

Nephropathogenic Infectious Bronchitis (NIB) is an important viral disease of poultry associated with severe renal damage and economic losses. The present study investigated the haematological alterations associated with naturally occurring NIB in broiler chickens from five confirmed flocks. Gross and histopathological examination revealed visceral uricosis, renal congestion, haemorrhage, necrosis, interstitial nephritis, glomerulonephritis, tubular degeneration and necrosis, and urate crystal deposition. Haematological analysis showed significantly reduced ($P < 0.05$) haemoglobin (Hb), total erythrocyte count (TEC), and total leukocyte count (TLC), indicating anaemia and leukopenia. Differential leukocyte count revealed significant heterophilia, lymphopenia, and reduced monocyte percentage, whereas eosinophil and basophil percentages did not differ significantly. These findings indicate that NIB is associated with characteristic haematological alterations that may serve as supportive indicators of disease in naturally infected broiler flocks.

Introduction

The emergence of NIB virus variants has raised serious concerns in the poultry industry due to their increased virulence and ability to cause severe renal damage,

resulting in high morbidity, mortality, and economic losses (Bayry *et al.*, 2005; Li *et al.*, 2022). Unlike the classical respiratory form, nephropathogenic strains exhibit a marked tropism for the kidneys, leading to nephritis, urate deposition, and impaired renal function

(Balasubramaniam *et al.*, 2012; Li *et al.*, 2022). Several researchers have documented characteristic gross and histopathological renal lesions associated with NIB, including visceral uricosis, nephritis, tubular degeneration and necrosis, and urate crystal deposition (El-Hamid *et al.*, 2016; Timurkaan *et al.*, 2023; Patel *et al.*, 2026). However, information on the haematological alterations associated with naturally occurring NIB under field conditions remains limited (Yohannes *et al.*, 2013; Yadav *et al.*, 2021). Therefore, the present study was undertaken to investigate the haematological changes associated with naturally occurring NIB in broiler chickens, in cases confirmed by characteristic gross and histopathological lesions.

Materials and Methods

Sample Collection

The study was carried out at the Department of Veterinary Pathology, College of Veterinary Science, AVFU, Khanapara, Guwahati-22. Carcasses of birds suspected of NIB were received from disease outbreak areas and organized commercial broiler farms of different districts of Assam for post-mortem examination. Representative tissue samples from affected kidneys were collected in 10% neutral buffered formalin for histopathological examination and were routinely processed and stained with Haematoxylin and Eosin (H&E) stain. Based on the gross and histopathological findings, five flocks were confirmed to be affected with NIB and were included in the study. Approximately 2 ml of blood was collected from the wing vein of clinically affected birds from the confirmed flocks in EDTA vials for routine haematological examination. Blood samples were also collected from apparently healthy birds and maintained in EDTA vials for comparison.

Haematological estimation

Total Erythrocyte Count (TEC) and Total Leukocyte Count (TLC)

A modified staining technique based on Tsai *et al.*, (2014), which was originally developed for red-eared slider turtles was used for TEC and TLC. The staining solution was formulated by mixing 0.1 ml of HiMedia® Gram's crystal violet (catalogue no. S012) solution with 9.9 ml of 0.45% sodium chloride solution. 10 µl of chicken blood and 740 µl of staining solution were

carefully mixed for cell counting, and the mixture was incubated for five minutes. After charging the hemacytometer with the stained sample, it was then allowed to stand for 10 minutes.

Red blood cells (RBCs) were counted in four small corner squares plus the centre square (1/25 mm²), while white blood cells (WBCs) were counted in four big corner squares (1 mm²). Cell counts were calculated using the following formulas:

1. **Total WBC count** = Total haematocytometer count ÷ 4 (large squares) ÷ 0.1 (depth of the chamber) x 75 (dilution rate).
2. **Total RBC count** = Total haemocytometer count x 5 (to reach 1-mm² area) ÷ 0.1 (depth of the chamber) x 75 (dilution rate).

Haemoglobin (Hb)

Sahli's apparatus was used to measure the Hb concentration using the acid hematin method, as detailed by Schalm *et al.*, (1975).

Differential leukocyte count (DLC)

DLC was performed in accordance with Schalm *et al.*, (1975).

Statistical Analysis

Student's t-test (Gosset, 1908) was used in SPSS software version 20 to statistically analyse the results from the haematological examination. P-values less than 0.05 (two-tailed) were deemed significant. The values for each parameter were presented as Mean ± SE.

Results and Discussion

Gross and Histopathology

Gross examination of NIB-affected birds revealed visceral uricosis with urate deposition on visceral organs (Figure 1) and severe renal involvement characterized by congestion, haemorrhage, necrosis (Figure 2), urate accumulation, and ureteral distension. Histopathologically (Figure 3), the kidneys showed interstitial nephritis, glomerulonephritis, tubular degeneration and necrosis, mononuclear cell infiltration, urate crystal deposition, and tubular epithelial

hyperplasia. Similar gross and microscopic lesions have been reported by [Munuswamy et al., \(2021\)](#), [Timurkaan et al., \(2023\)](#) and [Patel et al., \(2026\)](#).

Haematological studies

Four haematological parameters were examined in the blood samples: Haemoglobin (Hb), Total Erythrocyte Count (TEC), Total Leukocyte Count (TLC), and Differential Leukocyte Count (DLC). The mean values of these parameters along with their standard error (SE), for apparently healthy group and NIB-affected group has been presented in Table 1.

Haemoglobin, Total Erythrocyte Count and Total Leukocyte Count

The mean Hb and TEC values were significantly reduced ($P < 0.05$) in the NIB-affected group (8.98 ± 0.28 g/dl and $1.81 \pm 0.04 \times 10^6/\mu\text{l}$, respectively) than in the apparently healthy group (11.00 ± 0.23 g/dl and $2.29 \pm 0.05 \times 10^6/\mu\text{l}$, respectively), suggesting anaemia, which is in agreement with the findings of [Yadav et al., \(2021\)](#). Leukopenia was indicated by significantly reduced ($P < 0.05$) mean TLC values ($12.00 \pm 0.35 \times 10^3/\mu\text{l}$) in the NIB-affected group compared to the apparently healthy group ($16.96 \pm 0.49 \times 10^3/\mu\text{l}$). [Yohannes et al., \(2013\)](#)

and [Yadav et al., \(2021\)](#) reported similar findings.

Differential Leukocyte count

Heterophilia was indicated by a significantly higher ($P < 0.05$) mean percentage of heterophils in the NIB-affected group ($17.40 \pm 0.58\%$) than in the apparently healthy group ($10.90 \pm 0.48\%$), which is in agreement with the findings of [Yohannes et al., \(2013\)](#) and [Yadav et al., \(2021\)](#). The mean percentage of lymphocytes was significantly reduced ($P < 0.05$) in the NIB-affected group ($73.40 \pm 0.42\%$) compared to the apparently healthy group ($78.00 \pm 0.66\%$), indicating lymphopenia, which corroborates the findings of [Yohannes et al., \(2013\)](#) and [Yadav et al., \(2021\)](#). The mean percentage of monocytes was significantly reduced ($P < 0.05$) in the NIB-affected group ($8.50 \pm 0.45\%$) compared to the apparently healthy group ($10.20 \pm 0.55\%$); however, [Yohannes et al., \(2013\)](#) reported no significant difference, whereas [Yadav et al., \(2021\)](#) observed a significant increase in monocyte counts. There was no significant difference ($P > 0.05$) in the mean percentages of eosinophils and basophils between the NIB-affected group ($0.40 \pm 0.69\%$ and $0.30 \pm 0.15\%$, respectively) and the apparently healthy group ($0.50 \pm 0.16\%$ and $0.40 \pm 0.16\%$, respectively), which is consistent with the findings of [Yohannes et al., \(2013\)](#) and [Yadav et al., \(2021\)](#).




Figure.1 Carcasses showing deposition of urates on visceral surfaces




Figure.2 Kidneys showing areas of necrosis




Figure.3 Photomicrograph of kidney revealed urate crystal deposition in renal tubules, tubular necrosis and interstitial nephritis (H&E, 40X)

Table.1 Mean ± SE of different haematological parameters of NIB-affected birds and apparently healthy birds' group			
S.N.	Parameters	NIB-affected	Apparently healthy
1.	Hb (g/dl)	8.98* ± 0.28	11.00 ± 0.23
2.	TEC (10 ⁶ /μl)	1.81* ± 0.04	2.29 ± 0.05
3.	TLC (10 ³ /μl)	12.00* ± 0.35	16.96 ± 0.49
4.	DLC (%)		
a.	Heterophils	17.40* ± 0.58	10.90 ± 0.48
b.	Lymphocytes	73.40* ± 0.42	78.00 ± 0.66
c.	Monocytes	8.50* ± 0.45	10.20 ± 0.55
d.	Eosinophils	0.40 ± 0.69	0.50 ± 0.16
e.	Basophils	0.30 ± 0.15	0.40 ± 0.16

P<0.05 = *; P>0.05 = non-significant. Means bearing superscript indicate a statistically significant difference (P<0.05).

In conclusion, NIB in broiler chickens was characterized by visceral uricosis and severe renal lesions including interstitial nephritis, glomerulonephritis, tubular degeneration and necrosis, mononuclear cell infiltration, and urate crystal deposition.

Haematological examination revealed anaemia, leukopenia, heterophilia, lymphopenia, and a significant

reduction in monocyte percentage in affected birds.

These findings suggest that haematological evaluation may provide useful supportive information in the diagnosis and assessment of NIB under field conditions. In addition, the modified staining technique described by Tsai *et al.*, (2014), originally developed for red-eared slider turtles, was successfully adapted in the present

study for estimation of TEC and TLC in chickens, highlighting its potential utility for avian haematological investigations.

Acknowledgements

Authors are thankful to the Department of Veterinary Pathology, College of Veterinary Science, AVFU, Khanapara, Guwahati, India-781022 to carry out the part of this research work.

Author Contributions

Dr. Sakshi Patel: Conceptualization, Methodology, Investigation, Formal analysis, Data curation, Writing – original draft preparation, Writing – review & editing, Visualization. Dr. Biswajit Dutta: Supervision, Conceptualization, Validation, Methodology, Writing – review & editing. Dr. Pankaj Deka: Resources, Methodology, Investigation, Provision of experimental material (stain), Validation. Dr. S. A. Begum: Investigation, Resources, Validation, Supervision support. Dr. Rupam Dutta: Advisory support, Validation, Review of experimental design and manuscript. Dr. Abhijit Deka: Methodology support, Validation, Supervision. Dr. Suchand Doloi: Investigation, Data collection (sample collection support), Validation. Dr. Ajay Mendake: Writing – review & editing, Visualization support, Manuscript assistance.

Data Availability

The datasets generated during and/or analyzed during the current study are available from the corresponding author on reasonable request.

Declarations

Ethical Approval Not applicable.

Consent to Participate Not applicable.

Consent to Publish Not applicable.

Conflict of Interest The authors declare no competing interests.

References

Balasubramaniam, A., Balasubramaniam, G.A., Sivaseelan, S., Gopalakrishnamurthy, T.R., Dorairajan, N. and Dinakaran, A.M., 2012.

Pathotyping of infectious bronchitis virus isolated from broiler chickens showing nephropathy. *Indian Journal of Veterinary Pathology*, 36(1), pp.49-53.

Bayry, J., Goudar, M.S., Nighot, P.K., Kshirsagar, S.G., Ladman, B.S., Gelb Jr, J., Ghalsasi, G.R. and Kolte, G.N., 2005. Emergence of a nephropathogenic avian infectious bronchitis virus with a novel genotype in India. *Journal of clinical microbiology*, 43(2), pp.916-918. <https://doi.org/10.1128/jcm.43.2.916-918.2005>

El-Hamid, A., Hatem, S., Ellakany, H.F., Elkholia, S.Z., Elbestawy, A.R. and Diab, E.R., 2016. Incidence of Avian Nephritis and Infectious Bronchitis Viruses in Broilers in Egypt. *Alexandria Journal of Veterinary Sciences*, 51(2).

Li, N., Huang, C., Chen, W., Li, Z., Hu, G., Li, G., Liu, P., Hu, R., Zhuang, Y., Luo, J. and Gao, X., 2022. Nephropathogenic infectious bronchitis virus mediates kidney injury in chickens via the TLR7/NF- κ B signaling axis. *Frontiers in Cellular and Infection Microbiology*, 12, p.865283. <https://doi.org/10.3389/fcimb.2022.865283>

Munuswamy, P., Ramakrishnan, S., Latheef, S.K., Kappala, D., Mariappan, A.K., Kaore, M., Anbazhagan, K., Puvvala, B., Singh, K.P. and Dhama, K., 2021. First description of natural concomitant infection of avian nephritis virus and infectious bronchitis virus reveals exacerbated inflammatory response and renal damage in broiler chicks. *Microbial pathogenesis*, 154, p.104830. <https://doi.org/10.1016/j.micpath.2021.104830>

Patel, S., Dutta, B., Deka, P., Begum, S.A., Dutta, R., Buragohain, M., Dutta, A. and Phukan, G., 2026. Pathomorphological changes in the liver and kidney of broiler chickens infected with nephropathogenic infectious bronchitis virus. *International of Advanced Biochemistry Research*, 10(2), pp.378-381. <https://www.doi.org/10.33545/26174693.2026.v10.i2e.7607>

Schalm, O.W., Jaim, N.C. and Carroll, E.J., 1975. *Veterinary Haematology*. 3rd (ed.) Lea. *And Febiger, Philadelphia, USA*.

Timurkaan, N., Öngör, H., Kalender, H., Karabulut, B., Çöven, F., Çevik, A., Eröksüz, H. and Çetinkaya, B., 2023. Pathological and molecular findings of visceral gout caused by Israel variant 2 (IS/1494/06) genotype of infectious bronchitis virus in chickens. *Ankara Üniversitesi Veteriner*

- Fakültesi Dergisi*, 70(2), pp.149-156.
<https://doi.org/10.33988/auvfd.974221>
- Tsai, C.Y., Yu, J.F., Wang, Y.W., Fan, P.C., Cheng, T.Y. and Wang, L.C., 2014. An alternative staining method for counting red-eared slider turtle (*Trachemys scripta*) blood cells using crystal violet in cells diluted with 0.45% sodium chloride. *Journal of Veterinary Diagnostic Investigation*, 26(5), pp.610-615.
<https://doi.org/10.1177/1040638714542868>
- Yadav, J.V., Lakshman, M., Madhuri, D. and Kannaki, T.R., 2021. Hematological Alterations in the Colored Broiler Chicken Experimentally Infected with Nephro-Pathogenic Avian Infectious Bronchitis Virus. *Int. J. Curr. Microbiol. App. Sci*, 10(01), pp.1937-1946.
<https://doi.org/10.20546/ijcmas.2021.1001.225>
- Yohannes, T., Sharma, A.K., Singh, S.D. and Sumi, V., 2013. Experimental haematobiochemical alterations in broiler chickens fed with T-2 toxin and co-infected with IBV. *Open journal of veterinary medicine*, 3(05), pp.252-258.
<http://dx.doi.org/10.4236/ojvm.2013.35040>

How to cite this article:

Sakshi Patel, Biswajit Dutta, Pankaj Deka, S. A. Begum, Rupam Dutta, Abhijit Deka, Suchand Doloi and Ajay Mendake. 2026. Haematological Alterations Associated with Naturally Occurring Nephropathogenic Infectious Bronchitis in Broiler Chickens. *Int.J.Curr.Microbiol.App.Sci*. 15(6): 170-175.

doi: <https://doi.org/10.20546/ijcmas.2026.1506.017>