

Original Research Article

<http://dx.doi.org/10.20546/ijcmas.2016.506.090>

Floristic composition of medicinal plants of Nawada District, Bihar, India

S. M. Ateeque Ahmad¹, Baidyanath Kumar^{2*} and Nandji Kumar³

¹Research Scholar, PG. Department of Botany, Magadh University, Bodh Gaya, Bihar, India

²Visiting Professor, Department of Biotechnology, Patna Science College,
Patna, Patna University, Bihar, India

³Professor, PG. Department of Botany Magadh University, Bodh Gaya, Bihar, India

**Corresponding author*

ABSTRACT

The biodiversity, floristic composition and structure of medicinal plants in the fourteen blocks of Nawada District were studied. A total of 105 plant species belonging to 34 families, 75 genera and 5 life forms were recorded. Fabaceae, Moraceae, Meliaceae and Apocyanaceae were the overall diverse families (in terms of species richness) of the adult species, contributing 44.5% of all the species in the study. Trees were the most dominant life form (48.5%) followed by lianas (16.8%), herbs (10.9%), epiphytes (8.9%), shrubs (3.9 %) and the others (4.7%). Species richness among all life forms was highest in the Nda (90.5%) followed by Ap (87.6%), Pb (77%), Sda (73%), Ha (70.25%) and Gp (68.95%). Fabaceae, Moraceae and Meliaceae and Apocyanaceae were the most diverse families distributed in all the fourteen blocks of Nawada. The trees in all the forest types studied were generally tall. The difference in height of tree species could be partly explained by degradation in the form of logging of tall and big trees which has undoubtedly affected the vertical structure. Even though tree size (dbh) correlated with tree height in all the forest types, the relationship was stronger ($r = 0.741$ to 0.368 ; $p = 0.000- 0.002$). Thus, dbh of trees could be a better predictor of tree height. The forest reserve of Nawada looks floristically rich and structurally complex in the face of logging, farming activities and invasion in some parts of the forest. Thus, there is the need to curb the anthropogenic activities and plant invasion so as to protect the integrity of the forest including medicinal plants.

Keywords

Biodiversity,
Floristic
composition
and structure,
Medicinal plants
and Forest areas.

Article Info

Accepted:
25 May 2016
Available Online:
10 June 2016

Introduction

India has a rich herbal heritage which has been described in Ayurveda. It has 16 different agro climatic zones. A large number of plants are available in different bio-geographical zones of India (Gharpure, 2011; Ravi kumar, 2011). According to WHO, 20,000 plant species are used for medicine out of 2, 50,000 all over the world.

Of these eight hundred species are being used commercially. In India, the rural population uses about 8000 herbal plants for medicine. India having a rich traditional knowledge of herbs in the Vedas and ancient days remains marginalized in the fields of research on plants and its medicinal use. The country has a rich herbal heritage and

about 1, 22,600 hectares of land is under cultivation for medicinal plants.

Nawada district is one of the thirty-eight districts of Bihar State and Nawada town is the administrative headquarter of this district. It occupies an area of 2,494 square kilometers and is located at 24.88N 85.53E with an average elevation of 80 meters (262 feet). This district lies on both sides of the river Khuri in 24^o53'N and 85^o 33'E, and includes fourteen blocks. The climatic conditions of Nawada are somewhat dry and healthy. The winter, summer and rainy seasons are well marked. The overall rainfall is 1142.3mm with maximum i.e. around 92% of rainfall being received during monsoon months.

Modern medicines use many ingredients which have side effect on human organs. This has opened up new vistas of research in herbal medicines. Herbal medicines have no side effect in comparison to the synthetic drugs (Cragg, et al.,1996; Haslam, 2003). Important drugs have been analyzed from plants of Indian origin (Stermitz, et al., 1999; Osborn et al., 2000). *Curcuma longa* (turmeric), *Carica papaya* (Papaya) *Acacia catechu*, *Mentha* species, *Nicotiana tabacum* are some of the important plants grown by the farmers of this area. Farmers of this area have adopted the cultivation of *Mentha* species, Amla, Neem, Gurmae, Sarp Gandha etc. These plants have great use in traditional therapy by Ayurveda as well as modern medicine.

The fertile land of Nawada has the potential to grow many herbs of therapeutic use. People of this area utilize "Tilkor Patta" for eating. The Tilkor (*Momordica monodelpha*) stimulates insulin secretion from pancreas and therefore it is used as a herbal product for treatment of diabetes. Not only this, one more species Gurich is found

in this region which is also an effective medicine for diabetes. Amla, Jamun, *Aloe vera* and Gugal are also found in plenty in this area. These herbs have got extensive use in modern medicine.

The plants of medicinal importance are mostly wild whose cultivation is an essential requirement. In Nawada ignorance and lack of preservation knowledge has resulted in the extinction of many plants of medicinal importance.

With the development of synthetic drugs, plant products lost their significance. In the last few decades, however, there has been more interest in drugs obtained from vegetable sources than at any time in history because of the success with the antibiotics, and other natural drugs such as *Rauwolfia* (for the treatment of mental disorder), *Podophyllum* (a cathartic, as well as for curing cancerous tumors in mice), *Aloe* (a cathartic, as well as for the treatment of atomic radiation burn, *Veratrum* (hypertensive agent), (Peyote) (psychoactive drug) and sundry others. Saponins obtained from many members of *Dioscoreaceae* and *agavaceae*, can be converted into cortisone, into male hormone (testosterone) and female hormones (estrogens and progesterone). Saponins have many potential uses in the treatment of rheumatoid arthritis and the female hormones are used in contraceptive pills.

The study of drugs and drug plants has developed into modern pharmacognosy which deals with the knowledge of history, botany, preservation and commerce of crude drugs. Nature has provided a rich storehouse of herbal remedies to cure all mankind's ills.

The information on drugs and drug plants whose efficacy in medicine has been established is available in various authentic

books known as 'Pharmacopoeia' and the drugs included therein are described as 'official'. The most important of these pharmacopoeia are the United States Pharmacopoeia, British Pharmaceutical Codex, Indian Pharmaceutical Codex and National Formulary.

The medicinal value of drugs is due to the presence of certain substances such as alkaloids, glycosides, resins, volatile oils, gums, tannins, etc. Some of these are powerful poisons if administered indiscriminately, while others are dangerously habit-forming. Even the most dangerous drugs can be of value to human beings, if judiciously employed. The danger of self medication is serious and extensive. People who believe themselves ill or simply off-colour physically or mentally, often use wonder-working drugs (happy pills) to relieve themselves from the tensions of modern living. These tranquilizers have proved so effective that their use has increased amazingly and has now outstripped all other drugs with the exception of the antibiotics.

The active principles of plant drugs are commonly more concentrated in storage organs. Roots, seeds, bark and leaves are much represented in the *Materia Medica*, flowers are less commonly used, while woods and woody parts of herbaceous stems are usually relatively inert.

Floristic composition and studies of medicinal plants are essential in view of their value in understanding the extent of plant biodiversity [World Observation Monitoring Centre (WCMC), 1992]. Knowledge of floristic composition is also useful in identifying important elements of plant diversity, protecting threatened and economic species, and monitoring the state of reserves, among others (Tilman, 1988;

Ssegawa and Nkuutu, 2006). Thus, the study of floristic composition and structure becomes more imperative in the face of the ever increasing threat to ecosystem. Studies have shown that composition and structure of plants are influenced by a number of factors (Klinge et al., 1995; Haugaasen et al., 2003). Prominent among these factors are disturbances which are thought to be key aspects, and the cause of local species variation within an ecosystem based on their intensity, scale and frequency (Hill and Curran, 2003; Laidlaw et al., 2007). Disturbances can alter the successional pattern and subsequent composition, diversity, and structure of the plant (Doyle, 1981; Busing, 1995). Logging which has immediate and direct effects on composition and structure (Parthasarathy, 2001) also creates canopy openings which may cause regeneration problems, especially in exposed conditions where soils dry out rapidly and nutrient loss through run-off becomes common. Canopy openings readily support the growth of invasive weeds and other herbaceous plants which usually interfere with regeneration and impede recovery of trees and shrubs (Epp, 1987; Hawthorne, 1993 and 1994; Madoffe et al., 2006). Invasive weeds threaten biodiversity by displacing native species and disrupting community structure (Parker et al., 1999; Richardson et al., 2000; Sala et al., 2000; Stein et al., 2000). Soil water availability is also considered a key factor for the regeneration, survival and growth of seedling communities (Lieberman and Lieberman, 1984; Ceccon et al., 2002). Light conditions influence regeneration pathways strongly (Haugaasen et al., 2003) and ultimately affect the composition and structure of flora of an ecosystem. It has been reported that light limitation alone may prevent seedling survival regardless of other resource levels (Tilman, 1982). Flooding as a limiting factor influences seedling and

sapling species distribution, and establishment (Klinge et al., 1995; Wittmann and Junk, 2003), probably as a result of physiological stress from highly anoxic conditions, as well as physical flood disturbance (Haugaasen and Peres, 2006). Magadh zone of Bihar is generally believed to be floristically rich, containing many tropical timber species and medicinal plants as well. Carefully compiled and up-to date information on these plant resources is, however, lacking except some fragmentary report (Jyoti and Baidyanath, 2015). Therefore, an attempt has been made to study the floristic composition of medicinal plants of Nawada District (Bihar).

Methodology

The study was carried out in the fourteen blocks covering almost all the areas of Nawada District (Bihar). These include:

Nawada (Nda), Akbarpur (Ap), Warisaliganj (Wg), Pakribarawan (Pb), Sirdala (Sda), Rajauli (Rjl), Roh (Rh), Kawakol (Kal), Hisua (Ha), Nardiganj (Ndg), Narhat (Nht), Gobindpur (GP), Meskaur (Mkr) and Kashichak (Kac).

The medicinal plants were collected from May 2015 to April 2016 and identified and their dbh (diameter at breast height) measured with a diameter tape. The height of all plants was determined with clinometers. In each quadrat all trees (dbh \geq 10 cm) were examined for the presence of climbers (lianas with dbh \geq 2 cm and vines). Trees were also surveyed for epiphytes according to the method described by Addo-Fordjour et al. (in press). Trees (dbh \geq 10 cm) were classified into four groups based on their height; understorey (< 20 m), low canopy (20-30 m), upper canopy (30-40 m) and emergent (> 40 m) species. The percentage canopy cover of each plot was

determined by a spherical densitometer. At each plot four readings from the four cardinal directions were taken at four different points. The average of all readings for plots in each block (14 readings) was calculated and used as the percentage canopy cover of that forest type (Anning et al., 2008). The diameter of lianas was determined at 1.3 m from the rooting base (Addo-Fordjour et al., 2009). Each quadrat of 25 m x 25 m was subdivided into twenty-five 5 m x 5 m small quadrates and fifteen of these (accounting for 60% of the plot area) were randomly sampled for herbs and regeneration of the tree species. All herbs with (\leq 2 m high with dbh < 10 cm) were identified and counted. Identification was performed by a plant taxonomist aided by manuals and Floras (Hawthorne, 1990; Arbonnier, 2004; Poorter et al., 2004; Hawthorne and Jongkind, 2006). Identification of the species was confirmed at the KNUST, Kumasi and the Forestry Commission, Kumasi herbaria.

Data Analyses

Differences in plant densities between the forest types were tested using one-way ANOVA. The relationship between tree size (dbh) and height in the forest blocks was determined by Pearson correlation analysis. These analyses were performed using Minitab 15 software at a significance level of 5%. The diversity of plant species in the forest types was quantified using the Shannon-Wiener species diversity index (Gimaret-Carpentier et al., 1998; Blanc et al., 2000; Parthasarathy, 2001). The Jaccard's index of similarity (*I*) was calculated for each pair wise plot comparison (Blanc et al., 2000) and this was used to generate a dendrogram showing floristic similarities. The index is given by;

$$I = \frac{C}{U_x + U_y + C} \times 100$$

Where,

C = number of species common to both plot X and Y

U_x = number of species found only in plot X

U_y = number of species found only in plot y.

Epiphytes were excluded from all the above mentioned analyses since their densities could not be determined. Importance value index of the species was calculated as the sum of the species relative density, relative frequency and relative dominance (Kiruki and Njung'e, 2007; Addo-Fordjour et al., 2008).

Besides these areas, some indigenous plants of Nawada used as medicine are also identified, and their medicinal values have been summarized.

Results and Discussion

Important medicinal plants in Nawada collected in rural areas of fourteen blocks, their habit and the families they belong have been presented in Table-1

In Magadh area of Bihar, the major plants used in indigenous system of medicines have been presented in Table-2

Family dominance of medicinal plant species in the Nawada District has been presented in Table-3

A total of 105 adult plant species were identified in the fourteen blocks of Nawada. These belonged to 34 families, 75 genera and 5 life forms (Table 1; Figure1 & 2). Fabaceae, Moraceae, Meliaceae and

Apocyanaceae were the overall diverse families (in terms of species richness) of the adult species, contributing 44.5% of all the species in the study (Figure 2). Trees were the most dominant life form (48.5%) followed by lianas (16.8%), herbs (10.9%), epiphytes (8.9%), shrubs (3.9 %) and the others (4.7%).

Generally, species richness among all life forms was highest in the Nda(90.5%) followed by the Ap (87.6%), Pb(77%), Sda (73%), Ha (70.25%) and Gp (68.95%). Fabaceae, Moraceae and Meliaceae and Meliaceae were the most diverse families distributed in all the fourteen blocks of Nawada. The most important families in the Nawada were Fabaceae, Moraceae, Meliaceae and Apocyanaceae (Figure 2). In forest areas of Wg, Rjl, Ha, Ndg, Nht, Mkr and Kac, the medicinal plant species were in the range of 48 to 59.8%. In the forest area of other blocks the incidence of medicinal plant species were low in the range of 29.97% to 40.30 % (Table- 4). *C. odorata* was the most dominant species of herb in terms of number of individuals accounting for 69% of individual herbs in the Nda and 45% of all the herbs in the forest areas of fourteen blocks.

There were a total of 4167 individuals of woody species (excluding epiphytes) identified in the forest areas of fourteen blocks. Trees were more abundant (2916 individuals/ha) followed by lianas (1603 individuals/ha) and shrubs (248 individuals/ha) (Table - 5). Tree density was greatest in the Kac (290/ha) followed by Gp (287/ha), Mkr (285/ha), Ha (280/ha), Sda (250/ha), Rh (215/ha) and Rjl (214/hac). Density of Liana was maximum in Pb(126/ha) followed by Sda (124/ha), Nda (121/ha), and Wg and Nht (108/ha each). Other areas have low density of Liana. Similarly density of shrubs was maximum in

Ab(19/ha) followed by Kal and Kac(17/ha), Rjl (18/ha), Rh (16/ha) and Sda (15/ha). Others have low density of shrubs. Shannon-Wiener index was greater in the Ap ($H' = 3.80$) compared to Mkr ($H' = 3.70$) and Rjl, Gp, Pb and Kal ($H' = 23.50$ to 3.60). Others have H' value of less than 3.0 (Table 5). Density of plant species differed significantly between the forest types ($F=8.96$; $df = 2$; $p = 0.000$). In all, *C. mildbraedii* was by far the most abundant species accounting for an average 10% of stems in all the habitats whereas *T. scleroxylon* was the most dominant species in terms of basal area representing 25% on the average (Table- 5). The overall dominant species in terms of the species importance value (average) were *T. scleroxylon* (28.2) and *C. mildbraedii* (23.7). The commonest species were *C. mildbraedii* and *Alafia barteri* with the average occurrence of 6.8 and 5.2 % respectively. On forest type basis *C. mildbraedii*, *C. zenkeri*, *L. welwitschii*, *_ansiona altissima*, *N. papaverifera* and *T. scleroxylon* were the dominant species in all the forest areas of eighteen blocks (Table 3). *B. papyrifera*, *C. mildbraedii*, *N. papaverifera*, *P. africanum* and *S. oblonga* were the dominant species in terms of basal area representing 25% on the average (Table 3). The overall dominant species in terms of the species importance value (average) were *T. scleroxylon* (28.2) and *C. mildbraedii* (23.7). The commonest species were *C. mildbraedii* and *_afia barteri* with the average occurrence of 6.8 and 5.2 % respectively.

On forest type basis *C. mildbraedii*, *C. zenkeri*, *L. welwitschii*, *_ansiona altissima*, *N. papaverifera* and *T. scleroxylon* were the dominant species in the in all the areas selected for present investigation (Table 3). *B. papyrifera*, *C. mildbraedii*, *N. papaverifera*, *P. africanum*, *S. oblonga* and *T. scleroxylon* were the species that

dominated the woody flora. In terms of size, majority of the trees were of the lower diameter class (10-30 cm) (Table-5). The number of individual trees in the categories decreased with increasing size of the trees. Larger diameter trees (90- 110 and > 110 cm) were not found in any areas selected for present investigation. Mean basal area recorded was in the range of 32.6 ± 0.8 (Wg) to 64.7 ± 2.6 m²/ha (Mkr) (Table 5). In the same way, mean canopy cover and height were higher in the Rjl (81.3 ± 4.3) followed by Gp(80.5 ± 3.4), Rh (78.5 ± 3.0), Kal (76.6 ± 3.5), Kac (76.2 ± 1.6), Ha (72.5 ± 1.5) and Nht (68.5 ± 2.8). The canopy cover of other forest areas ranged between 35.5 ± 0.8 and 64.7 ± 2.6 . In the present investigation it was found that there was a significant positive relationship between tree size and height in all the forest types selected for present studies ($r^2 = 0.812$; $p = 0.000$, $r^2 = 0.741$; $p = 0.000$ and $r^2 = 0.362$; $p = 0.002$).

Studies on biodiversity, floristic composition and structure in the forests of fourteen blocks of Nawada are instrumental in the sustainability of forests since they play a major role in the conservation of medicinal plant species, and the management of forest ecosystems as a whole (Tilman, 1988; Ssegawa and Nkuutu, 2006). That notwithstanding, only a few studies (Hall and Swaine, 1981; Vordzogbe et al., 2005; Anning et al., 2008) in this regard have been conducted in forest areas. For this reason, the results of this study cannot be compared with a wide range of other similar studies in different blocks of Nawada. On the whole, floristic composition of the forest regions of Nawada was found to be lower than many other tropical areas. For instance, Vordzogbe et al. (2005) reported of much higher species richness (80 species/ha) in a moist semi-deciduous forest in Ghana. Even at a higher diameter cut-off point of ≥ 30 cm, Parthasarathy (2001) recorded 125

species in a tropical wet evergreen forest in Sengaltheri of the Western Ghats in India. On the other hand, Anning et al. (2008) recorded a much lower species richness (37 species/ha) in another disturbed semideciduous forest located in Ghana. On the basis of only trees (dbh \geq 10 cm) the number of species recorded in this study (28 species/ha) was relatively lower in comparison with similar studies in other tropical forests. For example, Campbell et al. (1986) and Riswan (1987) recorded 189 species/ha and 160 species/ha in Brazilian Amazon and Lempake Indonesia respectively. The cutting of mature trees for timber, farming, collection of fuel wood and other non-timber forest products, and their attendant invasion of some parts of the forest might have had effects on the species composition of the reserve (Terborgh, 1992; Odoom, 2005; Opoku, 2006). Trees constituted the predominant life form in all the forest types reminiscent of other studies (Vordzogbe et al., 2005; Anning et al., 2008). Although shrubs and herbs were generally less prominent in all the forest types, the diversity of herbs was greater in all the areas of fourteen blocks of Nawada. This might be an indicator of greater anthropogenic disturbances in the forest areas of Nawada (Mishra et al., 2008). The comparatively high proportion of *Ficus* spp. in the epiphytic flora is good for the health of the forest reserve since a lot of *Ficus* spp. are keystone species in many tropical forests (Lambert and Marshall, 1991). The dominance of *C. mildbraedii* and *T. scleroxylon* in the flora of the forest areas is considered as a major characteristic of semi-deciduous forests (Taylor, 1960; Hall and Swaine, 1981; Vordzogbe et al., 2005). Furthermore, the dominance of Fabaceae, Moraceae and Meliaceae and Apocyanaceae in some semi-deciduous forests has been reported (Vordzogbe et al., 2005; Anning et al., 2008). Thus, the semi-deciduousness of

the forest reserve in Nawada as well as the dominance of these species (*C. mildbraedii* and *T. scleroxylon*) (Hall and Swaine, 1981) has been maintained for more than two decades. The presence of some species such as *A. barteri*, *B. bespubescens*, *C. mildbraedii* and *G. simplicifolia* in all the fourteen blocks may indicate their wider range of ecological adaptation (Senbeta et al., 2005; Addo-Fordjour et al., 2008). Floristic composition did not vary much between the various forest types of Nwathathan in the other forest types which continue to be disturbed by human activities. This is buttressed by the density of plant species which was also significantly greater ($p = 0.000$) in the Nda compared to the other blocks. Logging and farming activities in all the eighteen blocks have affected plant diversity through the removal of plant species, creation of gaps, and consequently a reduction in the canopy cover of these forests. Both canopy gaps and farming activities Nawada have favoured the growth of invasive species among the native species, with *C. odorata* and *B. papyrifera* which have been identified as notorious invasive weeds being common phenomena. Though invasive species have been found to exert severe negative consequences on biodiversity, displacing native species and disrupting community structure (Ambika, 1996; Parker et al., 1999; Richardson et al., 2000; Sala et al., 2000; Stein et al., 2000; Madoffe et al., 2006). This could be attributed to the early stage of invasion in the Nawada, indicating that invasion is yet to have its full impact on the flora of the area. Therefore, there is still an opportunity for management intervention to control *B. papyrifera* and *C. odorata* so as to protect the biodiversity of the forest. Compared with other tropical forests, the forest areas of Nawada vertically presented a higher structural complexity

The trees in all the forest types studied were generally tall. The difference in height of tree species could be partly explained by degradation in the form of logging of tall and big trees which has undoubtedly affected the vertical structure. Even though tree size (dbh) correlated with tree height in all the forest types, the relationship was stronger ($r = 0.741$ to 0.368 ; $p = 0.000-0.002$). Thus, dbh of trees could be a better predictor of tree height. Age of trees has been found to influence the relationship between tree height and dbh with younger forests showing a stronger relationship than matured forests (Ryan and Yoder, 1997). Thus, the cutting down of matured trees was at least partly responsible for the better correlation between these parameters compared to the other forest types. Generally, tree density in relation to vertical structure of the forest, decreased with increasing height of canopy layers. The cutting down of large trees could explain the relatively lower mean basal area of woody species in these forest types.

The greatest diversity of saplings (H') in the DB could be attributed to higher sunlight intensity in the forest floor (following gap creation) that supported growth and

development of seedlings and saplings. Invasion of the forest areas by *B. papyrifera* and *C. odorata* partly contributed to the poor regeneration of this forest. Sapling and seedling diversity decreased with increasing cover of *B. papyrifera* and *C. odorata* until there were no seedlings or saplings of the native species in areas where the invasive species had formed monotypic stands. This is supported by the findings of Sharma and Raghubanshi (2006). *B. papyrifera* alone constituted 47 % of all the regenerating stems in the DIF, showing their dominance among the regenerating species. The continual dominance of these species may likely affect the regeneration of more native species. The devastating effects of *B. papyrifera* on the regeneration capacity of some native species have also been reported (CSIR, 2002). Seedling diversity was poor in the Nda but the relatively higher numbers of saplings and trees in the smaller diameter class may indicate the ability of the Nda to recruit more seedlings into the adult phase. This finding is supported by the work of Mishra et al. (2008) and Jyotu Jyotsna and Baidyanath Kumar (2015) in which a higher rate of conversion of saplings to trees was recorded in undisturbed forests.

Table.1 Important Medicinal plants of Nawada

Species	Family	Habit
<i>Acacia kamerunensis</i> Gand.	Fabaceae	Liana
<i>Acanthaceae</i> sp	Acanthaceae	Herb
<i>Acacia pentagona</i> (Schum. & Thonn.) Hooker f	Fabaceae	Liana
<i>Afromomum</i> sp	Zingiberaceae	Herb
<i>Azelia bella</i> Harms	Fabaceae	Herb
<i>Aidia genipiflora</i> (DC.) Dandy	Rubiaceae	Tree
<i>Alafia barberi</i> Oliv	Apocynaceae	Liana
<i>Albizia adianthifolia</i> (Schum.) W.F. Wight	Fabaceae	Tree
<i>Albizia glaberrima</i> (Schum. & Thonn.) Benth	Fabaceae	Tree
<i>Albizia zygia</i> (DC.) J.F. Macbr	Fabaceae	Tree
<i>Alstonia boonei</i> De Wild	Apocynaceae	Tree
<i>Amphimas pterocarpoides</i> Harms	Fabaceae	Tree
<i>Anchomanes difformis</i> (Blume) Engl.	Araceae	Herb

<i>Antiaris toxicaria</i> (Rumph ex Pers.) Leschen	Moraceae	Tree
<i>Antrocaryon micraster</i> A. Chev. & Guillaum	Anacardiaceae	Tree
<i>Baphia nitida</i> Lodd.	Fabaceae	Tree
<i>Baphia pubescens</i> Hook.f.	Fabaceae	Tree
<i>Blighia sapida</i> Kon.	Sapindaceae	Tree
<i>Blighia welwitschii</i> (Hiern) Radlk	Sapindaceae	Tree
<i>Bombax buonopozense</i> P.Beauv.	Bombacaceae	Tree
<i>Bridelia atroviridis</i> Müll.Arg.	Euphorbiaceae	Tree
<i>Bridelia grandis</i> Pierre ex Hutch.	Euphorbiaceae	Tree
<i>Broussonetia papyrifera</i> Vent.	Moraceae	Tree
<i>Bussea occidentalis</i> Hutch.	Fabaceae	Tree
<i>Calpocalyx brevibracteatus</i> Harms	Fabaceae	Tree
<i>Calycobolus africanus</i> (G.Don) Heine	Convolvulaceae	Tree
<i>Calypstrochilum emarginatum</i> Schltr.	Orchidaceae	Epiphyte
<i>Ceiba pentandra</i> (L.) Gaertn.	Bombacaceae	Tree
<i>Centrocoma pubescens</i> Benth.	Fabaceae	Herb
<i>Chromolaena odorata</i> (L.) King & Robinson	Asteraceae	Herb
<i>Chrysophyllum perpulchrum</i> Mildbr. Ex Hutch. & Dalziel	Sapotaceae	Tree
<i>Chrysophyllum</i> sp.	Sapotaceae	Tree
<i>Cissus</i> sp.	Vitaceae	Liana
<i>Combretum bipindense</i> Engl. & Diels	Combretaceae	Liana
<i>Combretum smeathmannii</i> G.Don	Combretaceae	Liana
<i>Combretum</i> sp.	Combretaceae	Liana
<i>Cordia millenii</i> Baker	Boraginaceae	Tree
<i>Cordia senegalensis</i> Juss.	Boraginaceae	Tree
<i>Corynanthe pachyceras</i> K.Schum.	Rubiaceae	Tree
<i>Rhaphidophora 814uperb814e</i> N.E.Br.	Araceae	Vine
<i>Dalbergia 814uperb814e</i> Benth.	Fabaceae	Liana
<i>Diospyros viridicans</i> Hiern	Ebenaceae	Tree
<i>Entandrophragma angolense</i> (Welw.) DC.	Meliaceae	Tree
<i>Ficus asperifolia</i> Miq.	Moraceae	Shrub
<i>Ficus 814uperb814e814814</i> Vahl	Moraceae	Shrub
<i>Ficus sur</i> Forssk.	Moraceae	Tree
<i>Ficus tessellata</i> Warb.	Moraceae	Epiphyte
<i>Ficus thonningii</i> Blume	Moraceae	Epiphyte
<i>Ficus trichopoda</i> Baker	Moraceae	Epiphyte
<i>Ficus 814uperb814e814</i> Vahl	Moraceae	Epiphyte
<i>Ficus vogelii</i> Miq.	Moraceae	Epiphyte
<i>Funtumia 814uperb814</i> (Preuss) Stapf	Apocynaceae	Tree
<i>Griffonia simplicifolia</i> (Vahl ex DC.) Baill.	Fabaceae	Liana
<i>Guarea cedrata</i> (A.Chev.) Pellegr.	Meliaceae	Tree
<i>Hippocratea</i> sp.	Celastraceae	Liana
<i>Hymenostegia afzelii</i> (Oliv.) Harms	Fabaceae	Tree
<i>Khaya anthotheca</i> (Welw.) C.DC.	Meliaceae	Tree
<i>Khaya grandifolia</i> C.DC.	Meliaceae	Tree
<i>Khaya ivorensis</i> A.Chev.	Meliaceae	Tree
<i>Lannea welwitschii</i> (Hiern) Engl.	Anacardiaceae	Tree
<i>Lecaniodiscus cupanioides</i> Planch. Ex Benth.	Sapindaceae	Tree

<i>Leptoderris</i> sp.	Fabaceae	Liana
<i>Lovoa trichilioides</i> Harms	Meliaceae	Tree
<i>Macaranga heudelotii</i> Baill.	Euphorbiaceae	Tree
<i>Mansonia altissima</i> (A.Chev.) A.Chev.	Sterculiaceae	Tree
<i>Marantocloa leucantha</i> (K.Schum.) MilneRedh	Marantaceae	Herb
<i>Microdesmis puberula</i> Hook.f.	Pandaceae	Tree
<i>Microsorium punctatum</i> (L.) Copel.	Polypodiaceae	Epiphyte
<i>Microsorium scolopendria</i> Copel	Polypodiaceae	Epiphyte
<i>Milicia 815uperb</i> (Welw.) C.C.Berg.	Moraceae	Tree
<i>Millettia chrysophylla</i> Dunn	Fabaceae	Liana
<i>Morinda lucida</i> Benth.	Rubiaceae	Tree
<i>Morus mesozygia</i> Stapf	Moraceae	Tree
<i>Motandra guineensis</i> (Thonn.) A.DC.	Apocynaceae	Liana
<i>Myrianthus arboreus</i> P.Beauv.	Cecropiaceae	Tree
<i>Olyra latifolia</i> L.	Poaceae	Herb
<i>Panicum maximum</i> Jacq.	Poaceae	Grass
<i>Parquetina nigrescens</i> (Afzelius) Bullock	Asclepiadaceae	Liana
<i>Pennisetum purpureum</i> K.Schum.	Poaceae	Grass
<i>Piptadeniastrum africanum</i> (Hook.f.) Brenan	Fabaceae	Tree
<i>Pisonia 815uperb815e</i> L.	Nyctaginaceae	Liana
<i>Pteris</i> sp.	Pteridaceae	Fern
<i>Pycnanthus angolensis</i> (Welw.) Warb.	Myristicaceae	Tree
<i>Ricinodendron heudelotii</i> (Baill.) Pierre ex Pax	Euphorbiaceae	Tree
<i>Atropa belladonna</i> L.	Solanaceae	Herb
<i>Digitalis</i> spp. Linn.	Scrophulariaceae	Herb
<i>Rauwolfia serpentine</i> Benth. Ex Kurtz.	Apocyanaceae	Undershrub
<i>Catheranthus roseus</i>	Apocyanaceae	Herb
<i>Rinorea oblongifolia</i> (C.H. Wright) Marquand ex Chipp	Violaceae	Tree
<i>Salacia elegans</i> Welw. Ex Oliv.	Celastraceae	Liana
<i>Salacia owabiensis</i> Hoyle	Celastraceae	Liana
<i>Salacia</i> sp.	Celastraceae	Liana
<i>Smilax kraussiana</i> Meisn.	Smilacaceae	Liana
<i>Sterculia oblonga</i> Mast.	Sterculiaceae	Tree
<i>Sterculia rhinopetala</i> K.Schum.	Sterculiaceae	Tree
<i>Sterculia tragacantha</i> Lindl.	Sterculiaceae	Tree
<i>Terminalia 815uperb</i> Engl. & Diels	Combretaceae	Tree
<i>Trichilia monadelpha</i> (Thonn.) J.J.de Wild	Meliaceae	Tree
<i>Trichilia priureana</i> A.Juss.	Meliaceae	Tree
<i>Trilepisium madagascariense</i> DC	Meliaceae	Tree
<i>Capsicum</i> sp	Solanaceae	Herb
<i>Abutilon indicum</i>	Malvaceae	Shrub
<i>Adhatoda vasica</i>	Acanthaceae	Shrub
<i>Caesalpinia bonducela</i>	Caesulpinaceae	Tree
<i>Cryptostegia grandiflora</i>	Asclepiadaceae	Shrub

Table.2 Major plants used as indigenous medicines in Magadh area

Species	Family	Used in Diseases
<i>Calotropis gigantea</i>	Asclepiadaceae	Liver, intestine spleen, piles
<i>Adhotoda vasica</i>	Acanthaceae	T.B., Cough, Hyperplasia
<i>Clitoria ternatia</i> (Aprajita)	Fabaceae	Swelling, Mental headache, Jaundice
<i>Achyranthus Aspera</i>	Nyctaginaceae	Appetite, Eyesight
<i>Nelumbian speciosum</i>	Nymphaeaceae	Skin diseases, piles
<i>Cucumis utilissimus</i>	Cucurbitaceae	Urinary disorder, Gall bladder, cholesterol
<i>Nerium odorum</i>	Apocyanaceae	Skin diseases, joint pain
<i>Strychros Nuxvomica</i>	Loganiaceae	Nervous system, Joint pain, Dysentery
<i>Cucumis melo</i>	Cucurbitaceae	Urinary disorder, kidney pain
<i>Daucus Carota</i>	Umbelliferae	Bile, nervous system, Asthma
<i>Leucas cephalotus</i>	Lemiaceae	Rheumatism, Cough, Anorexia
<i>Calendula officinalis</i>	Asteraceae	Asthma, Cough, tooth, Skin disease
<i>Aloe vera</i>	Liliaceae	Liver, spleen, Skin disease
<i>Ricinus Communis</i>	Euphorbiaceae	Heart pain, joint pain, piles
<i>Mentha spp.</i>	Lemiaceae	Urinary disorder, Worm killer
<i>Eclipta alba</i>	Asteraceae	Cough, Skin diseases, Liver, spleen
<i>Moringa pterygosperma</i>	Moringaceae	Anticancer, painkiller, ulcer, cough, liver, spleen, fever
<i>Aconitum napellus</i>	Ranunculaceae	Neuralgia and Rheumatism
<i>Cassia angustifolia</i>	Caesalpinaceae	Laxative used in habitual constipation
<i>Withania somnifera</i>	Solanaceae	Diuretic, Rheumatism, applied to ulcer carbuncle

Fig.1 Composition of medicinal plant species in the various life forms identified in Nawada




Table.3 Family dominance of medicinal plants

Family	Number of Species
Fabaceae	18
Moraceae	13
Euphorbiaceae	4
Apocyanaceae	6
Combretaceae	4
Sterculiaceae	4
Meliaceae	8
Celastraceae	4
Poaceae	3
Bombaceae	2
Solanaceae	2
Polypodiaceae	2
Rubiaceae	2
Marantaceae	1
Ppandaceae	1
Asclepiadaceae	2
Nyctaginaceae	1
Ebenaceae	1
Acanthaceae	2
Zingiberaceae	1
Rubiaceae	2
Araceae	2
Anacardiaceae	2
Sapindaceae	2
Boraginaceae	2
Cecropiaceae	1
Bombaceae	2
Convulvulaceae	1
Orchidaceae	1
Asteraceae	1
Sapotaceae	2
Vitaceae	1
Malvaceae	1
Caesalpinaceae	2
Asclepiadaceae	2

Table.4 Abundance and dominance of woody medicinal plant species in all the fourteen blocks of Nawada District

Species	Nda	Ap	Wg	Pb	Sda	Rjl	Rh	Kal	Ha	Ndg	Nht	GP	Mkr	Kac
<i>Acacia kamerunensis</i>	-	-	0.25	0.55	0.57	0.25	0.75	1.25	-	-	-	-	0.65	-
<i>Acacia pentagon</i>	-	-	0.35	0.35	0.65	0.26	0.55	0.15	0.25	0.65	-	-	0.35	-
<i>Acanthaceae sp</i>	0.67	0.75	-	-	-	-	-	-	0.55	-	-	-	0.65	-
<i>Afromomum sp</i>	-	-	-	-	-	-	-	-	-	-	0.55	0.65	0.35	-
<i>Afzelia bella</i>	-	-	-	-	-	0.57	-	-	-	-	-	-	0.65	-
<i>Aidia genipiflora</i>	-	-	-	-	-	0.58	-	-	-	-	-	-	0.45	-
<i>Alafia barteri</i>	-	-	-	-	-	0.55	-	-	-	-	-	-	0.65	-
<i>Albizia adianthifolia</i>	2.55	3.5	5.6	6.5	7.5	7.5	4.5	4.5	6.5	7.5	3.4	7.5	6.5	-
<i>Albizia zygia</i>	3.5	2.6	-	-	-	1.5	-	-	-	-	-	0.75	6.0	-
<i>Alstonia boonei</i>	-	-	-	-	-	5.5	-	-	-	-	-	-	-	-
<i>Amphimas pterocarpoides</i>	-	-	-	7.5	-	-	-	-	-	5.5	-	-	-	6.5
<i>Anchomanes difformis</i>	-	-	-	-	-	-	-	-	-	-	8.5	-	-	-
<i>Antiaris toxicaria</i>	2.5	-	-	-	-	-	4.6	-	-	-	-	-	-	-
<i>Antrocaryon micraster</i>	-	-	0.5	-	-	-	1.5	-	3.5	-	-	-	-	-
<i>Baphia nitida</i>	-	-	-	-	-	-	-	-	-	-	-	-	-	-
<i>Baphia pubescens</i>	0.55	-	-	-	-	-	-	-	-	0.25	-	-	-	-
<i>Blighia sapida</i>	5.5	-	4.5	-	-	-	6.4	-	-	-	5.5	-	-	-
<i>Blighia welwitschii</i>	-	-	-	3.6	-	-	-	-	4.5	-	-	-	-	2.5
<i>Bombax buonopozense</i>	7.5	-	-	-	8.5	9.4	-	-	-	-	-	-	6.5	-
<i>Bridelia atroviridis</i>	-	-	-	-	-	-	-	-	-	-	-	-	5.5	-
<i>Bridelia grandis</i>	-	-	-	-	-	-	-	-	-	-	-	-	6.0	-
<i>Broussonetia papyrifera</i>	-	2.5	-	-	-	-	-	3.5	-	-	-	2.6	-	-
<i>Bussea occidentalis</i>	-	5.0	-	-	-	-	-	3.0	-	-	-	2.5	-	-
<i>Calpocalyx brevibracteatus</i>	-	-	-	4.5	-	-	-	-	3.5	-	-	-	-	2.7
<i>Calycobolus africanus</i>	-	-	-	-	-	-	-	-	4.6	-	-	-	-	-
<i>Calyptrochilum emarginatum</i>	-	-	-	-	-	-	-	-	2.5	-	-	-	-	-
<i>Capsicum sp.</i>	5.5	2.4	3.5	4.7	1.5	1.5	1.5	2.7	2.5	2.6	2.0	3.5	5.5	4.5
<i>Ceiba pentandra</i>	7.6	7.0	-	-	-	-	7.0	-	-	5.0	6.5	-	4.5	3.6
<i>Centrocema pubescens</i>	-	-	-	-	-	-	-	-	5.0	-	-	-	6.5	-

<i>Chromolaena odorata</i>	4.5	4.0	3.5	2.7	3.5	-	-	-	5.0	-	-	-	6.5	-
<i>Chrysophyllum perpulchrum</i>	6.5	6.0	4.0	3.5	1.2	0.5	-	0.6	-	0.4	-	-	-	0.7
<i>Chrysophyllum sp.</i>	-	0.5	-	0.7	-	-	-	1.5	-	-	2.0	-	-	-
<i>Cissus sp.</i>	-	-	-	-	-	-	2.6	-	-	-	-	5.0	-	-
<i>Combretum bipindense</i>	4.0	-	-	4.7	6.5	-	-	-	-	3.8	-	-	-	2.7
<i>Combretum smeathmannii</i>	3.5	-	-	3.4	4.5	-	3.0	-	-	4.0	-	-	-	3.0
<i>Combretum sp</i>	-	-	-	3.0	-	-	-	-	-	-	-	-	-	2.5
<i>Cordia millenii</i>	-	-	-	-	-	-	-	-	-	-	4.5	-	-	-
<i>Cordia senegalensis</i>	-	-	-	-	-	-	-	-	-	-	5.0	-	-	-
<i>Corynanthe pachyceras</i>	-	0.57	-	-	-	-	-	-	-	-	0.55	-	-	-
<i>Rhaphidophora africana</i>	-	0.7	-	-	-	-	-	-	-	-	0.5	-	-	-
<i>Dalbergia hostilis</i>	7.5	6.5	5.0	4.7	4.6	3.5	2.8	3.6	6.5	6.5	4.5	4.5	1.0	0.5
<i>Diospyros viridicans</i>	-	-	-	1.5	-	-	-	-	-	-	-	-	1.7	-
<i>Entandrophragma angolense</i>	0.5	-	-	-	-	-	-	-	-	0.7	-	-	-	-
<i>Ficus asperifolia</i>	2.5	-	-	-	-	3.6	-	-	-	-	-	-	5.5	-
<i>Ficus exasperata</i>	1.5	-	-	-	-	2.4	-	-	-	-	-	-	2.0	-
<i>Ficus sur</i>	2.5	-	-	4.5	-	-	6.5	-	-	-	-	1.5	-	-
<i>Ficus tessellata</i>	2.5	-	-	3.5	-	-	5.0	-	-	-	-	1.7	-	-
<i>Ficus thonningii</i>	0.5	0.7	-	0.5	-	0.6	-	-	-	-	-	-	0.5	-
<i>Ficus trichopoda</i>	0.7	0.4	-	-	-	0.5	-	-	-	-	-	0.5	-	0.7
<i>Ficus umbellata</i>	2.5	-	-	-	-	3.5	-	-	-	4.5	-	-	-	-
<i>Ficus vogelii</i>	-	-	-	-	-	-	-	-	0.7	-	-	-	-	-
<i>Funtumia elastica</i>	-	-	-	-	1.5	-	-	-	-	-	-	-	2.6	-
<i>Griffonia simplicifolia</i>	-	3.5	-	-	-	-	-	0.5	-	-	-	-	-	0.4
<i>Guarea cedrata</i>	-	-	5.0	-	-	-	-	-	-	-	4.5	-	-	-
<i>Hippocratea sp.</i>	-	-	-	0.3	-	-	-	-	-	-	-	-	-	-
<i>Hymenostegia afzelii</i>	-	-	-	-	-	-	-	-	-	-	-	-	-	0.5
<i>Khaya anthotheca</i>	-	-	-	-	-	-	-	-	-	-	-	-	-	0.4
<i>Khaya grandifolia</i>	-	-	-	-	-	-	-	-	-	-	-	-	-	0.5
<i>Khaya ivorensis</i>	-	-	-	-	-	-	-	-	-	-	-	-	-	0.3
<i>Lannea welwitschii</i>	-	0.6	-	-	-	-	-	-	-	0.5	-	-	-	-
<i>Lecaniodiscus</i>	-	0.5	-	-	-	-	5.5	-	-	-	-	-	-	-

<i>cupanioides</i>														
<i>Leptoderris sp.</i>	2.5	-	-	-	3.0	-	-	-	-	-	3.5	-	-	-
<i>Lovoa trichilioides</i>	-	-	0.7	0.5	-	0.6	-	1.0	-	-	-	-	0.5	-
<i>Macaranga heudelotii</i>	2.5	-	-	-	-	-	3.0	-	-	-	-	-	-	-
<i>Mansonia altissima</i>	-	0.6	-	-	-	-	-	-	0.7	-	-	-	-	-
<i>Marantocloa leucantha</i>	-	-	4.0	-	-	-	-	5.2	-	-	-	3.6	-	-
<i>Microdesmis puberula</i>	-	-	-	0.7	-	-	-	-	-	0.5	-	-	-	0.4
<i>Microsorium punctatum</i>	-	-	-	0.3	-	-	-	-	-	0.5	-	-	-	0.5
<i>Microsorium scolopendria</i>	-	-	-	0.5	-	-	-	-	-	0.5	-	-	-	0.6
<i>Milicia excelsa</i>	-	-	-	0.7	-	-	-	-	-	0.5	-	-	-	0.7
<i>Millettia chrysophylla</i>	-	0.5	-	-	-	-	-	-	0.7	-	-	-	-	-
<i>Morinda lucida</i>	-	-	-	-	2.5	-	-	-	-	-	-	2.7	-	-
<i>Morus mesozygia</i>	-	-	2.5	-	-	-	-	2.0	-	-	-	-	-	-
<i>Motandra guineensis</i>	-	-	2.0	-	2.4	-	-	1.5	-	-	-	1.7	-	-
<i>Myrianthus arboreus</i>	-	-	2.6	-	2.5	-	-	2.0	-	-	-	2.5	-	-
<i>Olyra latifolia</i>	-	-	3.5	-	-	-	-	-	-	-	-	-	-	-
<i>Panicum maximum</i>	-	2.7	-	-	4.0	-	-	-	-	1.8	-	-	-	-
<i>Parquetina nigrescens</i>	-	-	-	-	-	0.8	-	-	-	-	-	-	0.6	-
<i>Pennisetum purpureum</i>	-	-	-	-	-	0.7	-	-	-	-	-	-	0.5	-
<i>Piptadeniastrum africanum</i>	-	-	-	-	-	0.7	-	-	-	-	-	-	0.5	-
<i>Pisonia aculeata</i>	1.5	-	-	-	-	-	-	1.7	-	-	-	-	-	-
<i>Pteris sp.</i>	6.5	5.0	6.5	6.0	4.0	2.5	4.0	2.5	3.5	3.0	2.7	2.0	3.5	1.5
<i>Pycnanthus angolensis</i>	3.5	-	-	-	5.0	-	-	-	-	-	-	2.5	-	-
<i>Ricinodendron heudelotii</i>	1.5	-	-	0.7	-	-	-	-	0.5	-	-	-	0.5	-
<i>Atropa belladonna</i>	3.2	-	-	3.5	-	-	4.0	-	-	-	-	3.5	-	-
<i>Digitalis spp</i>	-	4.0	3.5	-	-	7.0	-	-	6.0	-	-	-	5.0	-
<i>Rauwolfia serpentine</i>	-	-	-	-	5.5	-	-	-	-	-	6.5	-	-	6.0
<i>Catheranthus roseus</i>	2.5	2.0	1.0	1.5	2.7	3.5	3.7	4.5	3.7	3.5	3.0	2.5	2.0	3.6
<i>Rinorea oblongifolia</i>	0.5	-	-	-	0.6	-	-	-	0.5	-	-	0.4	-	-

<i>Salacia elegans</i>	1.4	-	-	1.5	-	-	-	-	-	-	1.4	-	-	-
<i>Salacia owabiensis</i>	1.6	-	-	1.7	-	-	2.5	-	-	-	2.0	-	-	-
<i>Salacia sp.</i>	1.5	-	-	2.0	-	-	3.0	-	-	-	1.7	-	-	-
<i>Smilax kraussiana</i>	1.7	-	-	2.5	-	-	3.0	-	-	-	4.7	-	-	-
<i>Sterculia oblonga</i>	3.6	-	3.5	-	2.7	-	-	2.5	-	3.0	-	-	-	4.0
<i>Sterculia rhinopetala</i>	2.3	-	3.6	-	2.5	-	-	-	-	-	-	-	-	-
<i>Sterculia tragacantha</i>	-	-	-	-	4.5	-	-	-	-	-	-	-	-	2.7
<i>Terminalia superba</i>	1.5	3.5	-	5.0	-	-	-	-	-	-	-	1.4	-	-
<i>Trichilia monadelpha</i>	-	-	-	-	-	0.5	-	-	-	-	0.7	-	-	-
<i>Trichilia prieureana</i>	-	-	-	-	-	1.5	-	-	-	-	2.0	-	-	-
<i>Trilepisium madagascariense</i>	-	-	-	-	-	2.0	-	-	-	-	1.7	-	-	-
<i>Abutilon indicum</i>		1.3					0.5				1.5	0.5		
<i>Adhatoda vasica</i>		1.5					1.6				1.8	1.6		
<i>Caesalpinia bonducela</i>	1.7		2.5		2.6		1.4	1.6		1.5	0.5		0.5	
<i>Cryptostegia grandiflora</i>				2.0		2.5					1.7	1.6	2.0	1.4

Fig.2 Family dominance of medicinal plant species in Nawada District




Table.5 Summary characteristics of the Floristic composition and structure of the medicinal plants in fourteen blocks of Nawada

Characteristics	Nda	Ap	Wg	Pb	Sda	Rjl	Rh	Kal	Ha	Ndg	Nht	Gp	Mkr	Kac
Number of tree individuals (density/ha)	85	95	107	154	250	214	215	150	280	230	107	114	217	285
Number of individual shrubs (density/ha)	8	12	10	13	15	18	16	17	10	14	17	13	9	11
Density of Liana/ha	121	105	108	126	124	60	58	69	52	104	85	108	75	85
Shannon-Wiener Index	2.5	2.6	2.9	3.5	3.4	3.6	2.8	3.5	3.7	2.8	2.9	3.1	3.6	3.4
Shannon-Wiener Index for sampling	1.30	1.34	1.35	2.70	2.50	1.45	2.50	1.36	1.35	1.34	2.68	2.36	1.85	1.80
Mean canopy height (m)	35.5±3.5	36.6±3.2	29.8±3.4	32.5±2.5	34.40±2.6	28.30±1.58	27.50±1.75	26.35±2.50	27.25±3.12	32.15±1.78	30.26±2.15	29.10±2.16	27.35±1.85	35.15±1.26
Mean canopy cover (%)	54.3±4.5	60.5±2.5	61.5±2.6	64.5±2.4	71.2±4.2	81.3±4.3	78.5±3.5	76.6±3.5	72.5±1.5	68.5±5.0	56.2±2.6	58.5±3.6	59.5±4.5	80.5±3.4
Mean basal area (m ² /ha)	52.5±3.2	35.5±0.8	32.6±0.8	45.2±1.2	43.5±3.2	55.6±3.5	60.4±2.6	58.5±0.8	53.5±2.5	54.5±4.5	38.6±3.2	37.5±0.9	63.5±2.5	64.7±2.6

In conclusion, human disturbances have influenced the biodiversity, floristic composition of the district Nawada to some extent. Invasion on the other hand did not have much effect on floristic composition of the Nawada due to the early stage of invasion. Thus, proper management intervention is required to mitigate the impact of *B. papyrifera* and *C. odorata* before it gets out of control. Logging affected the structural complexity of the forest reserve through the removal of large and tall trees as well as gap creation. The forest regions of Nawada (Magadh area) had better capacity of recruiting saplings into the adult stage. Invasion of the DIF by *B. papyrifera* and *C. odorata* affected regeneration of native species. The forest reserve of Magadh looks floristically rich and structurally complex in the face of logging, farming activities and invasion in some parts of the forest. Thus, there is the need to curb the anthropogenic activities and plant invasion so as to protect the integrity of the forest including medicinal plants.

Acknowledgements

The first author is grateful to Dr. Baidyanath Kumar, Visiting Professor, Department of Biotechnology, Patna Science College, Patna for providing necessary suggestion and for helping in identification of plant species.

References

- Addo-Fordjour P, Anning AK, Addo MG, Osei MF (in press). Composition and distribution of vascular epiphytes in a tropical semideciduous forest, Ghana. *Afr. J. Ecol.*
- Addo-Fordjour P, Anning AK, Atakora EA, Agyei PS (2008). Diversity and distribution of climbing plants in a semi-deciduous rain forest, KNUST Botanic Garden, Ghana. *Int. J. Bot.* 4: 186-195.
- Addo-Fordjour P, Anning AK, Larbi JA, Akyeampong S (2009). Liana species richness, abundance and relationship with trees in the Bobiri forest reserve, Ghana: impact of management systems. *For. Ecol. Manage.* 257:1822-1828.
- Ambika SR (1996). Ecological adaptations of *Chromolaena odorata* (L.) King & Robinson. In: Proceedings of the Fourth International Workshop on Biological Control and Management of *Chromolaena odorata*, Bangalore, India, pp. 1–10.
- Anning AK, Akyeampong S, Addo-Fordjour P, Anti KK, Kwarteng A, Tettey YF (2008). Floristic composition and vegetation structure of the KNUST Botanic Garden, Kumasi, Ghana. *JUST* 28: 103-116.
- Arbonnier M (2004). Trees, shrubs and lianas of West African dry zones. CIRAD, MARGRAF Publishers.
- Blanc L, Maury-Lechon G, Pascal J-P (2000). Structure, floristic composition and natural regeneration in the forests of Cat Tien National Park, Vietnam: an analysis of the successional trends. *J. Biogeogr.* 27: 141–157.
- Busing RT (1995). Disturbance and the population dynamics of *Liriodendron tulipifera*: simulations with a spatial model of forest
- Campbell DG, Douglas CD, Prance GT, Maciel UN (1986). Quantitative ecological inventory of terra firme and varzea tropical forest on the Rio Xingu, Brazilian Amazon. *Brittonia* 38: 369-393.
- Ceccon E, Omstead I, Vázquez-Yanes C, Campo-Alves J (2002). Vegetation and soil properties in two tropical dry forests of differing regeneration status

- in Yucatán. *Agrociencia* 36: 621-631.
- Cragg, G. M DJ Newman, KM Snader: *Gen Pharmacol* 1996, 27(4): 713-22
- CSIR (2002). *Country Report on Invasive Alien Species in Ghana: Removing Barriers to Invasive Plant Management in Africa*. Council for Sci. and Ind. Res., Accra, Ghana, pp 49–57.
- Doyle TW (1981). The role of disturbance in the gap dy-namics of a montane rain forest: an application of a tropical forest succession model. In West DC, Shugart HH, Botkin DB (eds) *Forest succession: concepts and application*. Springer-Verlag, Berlin, Germany, pp. 56- 73.
- Epp GA (1987). The seed bank of *Eupatorium odoratum* along a successional gradient in a tropical rain forest in Ghana. *J. Trop. Ecol.* 3: 139-149.
- Gharpure, Y. H. *Chemical News*, June 2011, 22
- Gimaret-Carpentier C, Pelissier R, Pascal J-P, Houillier F (1998). Sampling tree species diversity in a dense moist evergreen forest with regard to its structural heterogeneity. *J. Veg. Sci.* 9: 161–172.
- Hall JB and Swaine MD (1988). Distribution and ecology of vascular plants in a tropical rain forest. Dr W. Junk Publishers, The Hague. succession. *J. Ecol.* 83:45-53.
- Haslam, E. *Nat Drod Rep* 2003 Feb; 20(1): 79-110
- Haugaasen T, Barlow J, Peres CA (2003). Surface wildfires in central Amazonia: Short-term impact on forest structure and carbon loss. *For. Ecol. Manage.* 179: 321-331.
- Haugaasen T, Peres CA (2006). Floristic, edaphic and structural characteristics of flooded and unflooded forests in the lower Rio Purús region of central Amazonia, Brazil. *Acta Amazon.* 36: 25–36.
- Hawthorne WD (1990). *Field guide to the forest trees of Ghana*. Chatham: Natural Resources Institute, for the Overseas Development Administration, London. Ghana Forestry Series 1.
- Hawthorne WD (1993). *Forest regeneration after logging: findings of a study in the Bia South Game Production Reserve, Ghana*. ODA Forestry Series. Natural Resources Institute, Chatham.
- Hawthorne WD (1994). *Fire damage and forest regeneration in Ghana*. ODA Forestry Series. Natural Resources Institute, Chatham.
- Hawthorne WD, Jongkind C (2006). *Woody plants of western African forests: a guide to the forest trees, shrubs and lianes from Senegal to Ghana*. Royal Botanic Gardens, Kew, UK.
- Hill JL, Curran PJ (2003). Area, shape and isolation of tropical forest fragments: effects on tree species diversity and implications for conservation forest area. *J. Biogeogr.* 30: 1391–1403.
- Jyoti Jyotsna and Baidyanath Kumar (2015): Biodiversity and Floristic composition of medicinal plants of Darbhanga, Bihar, India. *International Journal of Current Microbiology and Applied Science*, 4 (12), 263-283
- Klinge H, Adis J and Worbes M (1995). The vegetation of a seasonal várzea in the lower Solimões river, Brazilian Amazon. *Acta Amazonica* 25: 201-220.
- Laidlaw M, Kitching R, Goodall K, Small A, Stork N (2007). Temporal and spatial variation in an Australian tropical rainforest. *Austral Ecol.* 32: 10–20.
- Lambert FR, Marshall AG (1991). Keystone characteristics of birddispersed *Ficus* in a Malaysian Lowland Rain Forest.

- J. Ecol. 79:793- 809.
- Lieberman D, Lieberman M (1984). The causes and consequences of synchronous flushing in a dry tropical forest. *Biotropica* 16: 193-201.
- Madoffe S, Hertel GD, Rodgers P, O'Connell B, Killenga R (2006). Monitoring the health of selected eastern arc forests in Tanzania. *Afr. J. Ecol.* 44: 171-177.
- Mishra RK, Upadhyay VP, Mohanty RC (2008). Vegetation ecology of the Similipal Biosphere Reserve, Orissa, India. *Appl. Ecol. Env. Res.* 6: 89-99.
- Odoom FK (2005). Chainsawing in the Natural Forests of Ghana: An assessment of the socio-economic impacts of this practice, Forest Harvesting Case-Study 21 Food and Agriculture Organization of the United Nations, Rome.
- Opoku K (2006). Forest governance in Ghana: An NGO perspective. A report produced by Forest Watch Ghana for FERN. Zuidam Uithof, Utrecht, Netherlands.
- Parker IM, Simberloff D, Lonsdale WM, Goodell K, Wonham M (1999). Impact: toward a framework for understanding the ecological effects of invaders. *Biol. Invasions* 1: 3-19.
- Parthasarathy N (2001). Changes in forest composition and structure in three sites of tropical evergreen forest around Sengaltheri, Western Ghats. *Curr. Sci.*, 80: 389-393.
- Poorter L, Bongers F, Kouamé FN, Hawthorne WD (Eds) (2004). Biodiversity of West African forests: an ecological atlas of woody plant species. CABI publishers.
- Richardson DM, Pyešek P, Rejmánek M, Barbour MG, Panetta FD, West CJ (2000). Naturalization and invasion of alien plants: concepts and definitions. *Divers. Distrib.* 6: 93-107.
- Ravi kumar et. al., *Indian J Exp Biol* 2011, 49, 455
- Riswan S (1987). Structure and floristic composition of a mixed dipterocarp forest at Lempake, East Kalimantan. In: Kostermans, A.G.J.H. (Ed.), *Proceedings of the Third Round Table Conference on Dipterocarps* (16-20 April 1985, Mulawarman University, Samarinda, East Kalimantan, Indonesia), pp 435-457.
- Ryan MG, Yoder BJ (1997). Hydraulic Limits to Tree Height and Tree Growth. *Bio. Sci.*, 47: 235-242.
- Sala OE, Chapin FS, Armesto JJ, Berlow E, Bloomfield J (2000). Global biodiversity scenarios for the year 2100. *Sci.* 287: 1770-1774.
- Senbeta F, Schmitt C, Denich M, Demissew S, Vlek PLG, Preisinger H, Woldemariam T, Teketay D (2005). The diversity and distribution of lianas in the Afromontane rain forests of Ethiopia. *Divers. Distrib.* 11: 443-452.
- Sharma GP, Raghubanshi AS (2006). Tree population structure, regeneration and expected future composition at different levels of *Lantana camara* L. invasion in the Vindhyan tropical dry deciduous forest of India. *Lyonia* 11: 27-39.
- Ssegawa P, Nkuutu DN (2006). Diversity of vascular plants on Ssesse Islands in Lake Victoria, Central Uganda. *Afr. J. Ecol.* 44: 22-29.
- Stein B, Kutner LS, Adams JS (2000). *Precious Heritage: The Status of Biodiversity in the United States*. Oxford: Oxford University Press, pp 399.
- Swaine MD, Hawthorne WD, Bongers F, Toledo Aceves M (2005). Climbing plants in Ghanaian forests. In: Bongers F, Parren MPE, Traore´ D (Eds) *Forest climbing plants of West Africa:*

- diversity, ecology and management. CAB International, Wallingford, Oxfordshire, UK, pp. 93–108.
- Taylor CJ (1960). Synecology and silviculture in Ghana. Thomas Nelson and Sons, London, UK.
- Terborgh J (1992). Maintenance of diversity in tropical forests. *Biotropica* 24: 253–292.
- Tilman D (1982). Resource Competition and Community Structure. Princeton University Press, NJ.
- Tilman D (1988). Plant strategies and the dynamics and structure of plant communities. Princeton University Press. Princeton, New Jersey.
- Vorddzobe VV, Attuquayefio DK, Gbogbo F (2005). The flora and mammals of the moist semi-deciduous forest zone in the Sefwi-Wiawso District of the Western Region, Ghana. *W. Afr. J. Appl. Ecol.* 8: 49-64.
- Wittmann F, Junk WJ (2003). Sapling communities in Amazonian whitewater forests. *J. Biogeogr.* 30: 1533-1544.
- WCMC (1992). World Conservation Monitoring Centre. Global Biodiversity: Status of Earth's Living Resources. Chapman and Hall, London, UK.

How to cite this article:

Ateeque Ahmad, S. M., Baidyanath Kumar and Nandji Kumar. 2016. Floristic composition of medicinal plants of Nawada District, Bihar, India. *Int.J.Curr.Microbiol.App.Sci.* 5(6): 806-826. doi: <http://dx.doi.org/10.20546/ijcmas.2016.506.089>