

## Original Research Article

<https://doi.org/10.20546/ijcmas.2017.607.100>

## Response of Neem Cake Rhizobium and Inorganic Fertilizer on Soil Health Growth and Yield of Green gram (*Vigna radiata* L.) var. Samrat

Rajesh and Arun Alfred David\*

Department of Soil Science and Agricultural Chemistry, Sam Higginbottom University of Agriculture, Technology and Sciences, Allahabad, 211007(U.P.), India

\*Corresponding author

### ABSTRACT

A study was conducted during *kharif* season 2016-17 to study the “Response of Neem cake *Rhizobium* and Inorganic Fertilizer on Soil Health Growth and Yield of Green gram (*Vigna radiata* L.) var. Samrat” on crop research farm department of Soil Science and Agricultural Chemistry, Naini Agricultural Institute, SHUATS, Allahabad. The design applied for statistical analysis was carried out with 3x2x2 factorial randomized block design having three factors with three levels of NPK @ 0, 50, and 100 % ha<sup>-1</sup>, two level of Neem cake and *Rhizobium* @ 0, 100% respectively. The best treatment was T<sub>11</sub>-I<sub>2</sub>+N<sub>1</sub>+R<sub>1</sub> (@ 100%N P K ha<sup>-1</sup>+ 100% Neem cake and *Rhizobium*) that showed the significant increase on enrichment of soil fertility status. The textural classes’ names are given to soils based upon the relative proportion of each of the three soil separates – sand, silt and clay. Sandy loam in which sand is 61.21%, silt 25.36% and clay 13.46%. The combination of N P K, Neem cake and *Rhizobium* was show resulted in a slight decrease in particle density (2.33 Mgm<sup>-3</sup>), bulk density (1.11Mgm<sup>-3</sup>), pH 6.97 and EC 0.28 dSm<sup>-1</sup> in post-harvest soil. The maximum increased values were recorded of O.C (0.74 %), pore space (59.58%), water holding capacity (56.62%) and available macro nutrient as Nitrogen (314.84 kg ha<sup>-1</sup>), Phosphorus (33.45 kg ha<sup>-1</sup>) and Potassium (181.05kg ha<sup>-1</sup>) with treatment combination T<sub>11</sub>-[I<sub>2</sub>+ N<sub>1</sub>+ R<sub>1</sub>] (@ 100% N P K ha<sup>-1</sup>+ 100% Neem cake and *Rhizobium*). The maximum cost benefit ratio was recorded 1:2.73 and maximum net return Rs. 58651 per hectare and maximum yield 13.58q ha<sup>-1</sup> in treatment combination T<sub>11</sub>-[I<sub>2</sub>+N<sub>1</sub>+R<sub>1</sub>@ 100% N P K ha<sup>-1</sup>+ 100% Neem cake and *Rhizobium*].

#### Keywords

Soil nutrients,  
Yield attributes,  
N P&K, Neem  
cake, *Rhizobium*  
and Green gram.

#### Article Info

Accepted:  
14 June 2017  
Available Online:  
10 July 2017

### Introduction

In India, production of pulses is around 13.5-15 million tonnes during the last decade, while annual domestic demand is 18-19 million tons. The yield of pulses has remained virtually stagnant for the last 40 year (539 kg/ha in 1961 to 544 kg/ha in 2001 to 651 kg/ha in 2013-14).

India is short of supply by 2 to 3 million tonnes annually. (Agropedia, 2014-15). Green gram

[*Vigna radiata* L.] (2n= 2x= 22) is third important pulse after chickpea and Pigeon pea. It is a self-pollinated crop and is an important grain legume of the tropical area. Greengram is also called mung, moong, mungo, greengram, goldengram, chicksaw pea and Oregon pea. It belongs to the Family Fabaceae and sub Family Papilionaceae. The center of origin is India (Karpechenko, 1925). Majority of Indian population is vegetarian,

pulses are cheap and best source of protein for Indian diet. It contains 20-25 percent protein, which is more than two times of cereals. India is importing about 3 million tonnes and the future demand of pulses by 2017 will be 28.0 million tonnes (Anonymous, (2011)).

In India, it is cultivated in Maharashtra, Andhra Pradesh, Rajasthan, Orissa, Karnataka and Uttar Pradesh. Green gram contributes 18.07 % of total pulses area and 11.48 % of total pulses production in India.

Area, production and productivity of green gram in India are 34.4 lakh ha, 14 lakh tonnes and 406.98 kg/ha respectively (lipr.res.in, 2014-15).

Green gram is an excellent source of protein. It is rich in lysine and deficient in methionine and legume, which is free from anti-nutritional factor. The range for protein content is fairly wide (20- 24% per 100g of dry grain).

Temperature of 28°C to 33°C is optimum for seed germination and plant growth. Green gram plants are erect with few branches carrying pods borne in cluster and near the top of the plant. Pods contain 8- 15 green seeds (Prasad, 2009).

The green gram (*Vigna radiata* L. Wilczek) is one of the important pulse crop because of its adaptation to short growth duration, low water requirement, soil fertility and is favoured for consumption due to its easy digestibility and low production of flatulence (Shil and Bandopadhyaya, 2007).

### **Nitrogen**

Green gram is capable of fixing atmospheric nitrogen, it responds to small quantity of nitrogenous fertilizers applied as starter dose. Application of 15-20 Kg Nha<sup>-1</sup> has been found optimum to get better response.

Application of higher dose of nitrogen may reduce nodule number and nodule growth and thus adversely affect the nitrogen fixation capacity.

### **Phosphorus**

Phosphorus plays a vital role in photosynthesis, respiration, energy storage, cell elongation and improves the quality of crops. Deficient plants may have thin, erect and spindly stems and leaves turn into bluish-green colour. Phosphorus is an essential constituent of majority of enzymes, which are of great importance in the transformation of energy, in carbohydrate metabolism, in fat metabolism and also in respiration of plants. It enhances the activity of *Rhizobium* and increased the formation of root nodules. Thus, it helps in fixing more of atmosphere nitrogen in root nodules (Rajveer *et al.*, 2016).

### **Neem cake**

Neem cake is a potential source of organic manure. Neem has demonstrated considerable potential as a fertilizer. Our Neem cake also reduce alkalinity in soil, as it produce organic acid on decomposition, being totally natural, the Neem cake we offer hence ensure fertility of the soil, it also improve the organic matter contain of the soil, helping improvement in soil texture, water holding capacity, soil aeration for better root development.

The addition of Neem cake also positively affected the available soil organic carbon, N P K and Mn content of soil resulting better growth and grain yield of mung bean (*Vigna radiata* L.) besides suppressing soil borne pathogens (Murugan, 2011). The composition of Neem cake is 5.2 % N, 1.0 % P, 1.4 % K. Neem cake act as a nitrogen inhibitor means reduce the nitrification. It supplies the available nitrogen for a long time in the soil (Katyayan, 2012).

## **Rhizobium**

The use of biofertilizers are more eco-friendly in nature. They can play a significant role in fixing atmospheric nitrogen; biofertilizers enrich soil fertility and improves soil fertility. Of these biofertilizers, *Rhizobium* inoculants specific for different leguminous crop is the most important in India. The largest contribution of biological nitrogen fixation to agriculture is derived from the symbiosis between legumes and *Rhizobium* species (Meena *et al.*, 2016).

## **Materials and Methods**

A field experiment was conducted on research farm of department of Soil Science and Agricultural Chemistry SHUATS Allahabad, (U.P.) India. The soil of experimental area falls in order Inception and the experimental field is Alluvial in nature. The design applied for statistical analysis was carried out with 3x2x2 factorial randomized block design having three factors with three levels of N P K @ 0, 50, and 100% ha<sup>-1</sup>, two levels of Neem cake @ 0 and 100%ha<sup>-1</sup> and two level of *Rhizobium* @ 0 and 100% ha<sup>-1</sup> respectively (Table 1).

## **Experimental site**

The experiment was conducted at research Farm of Soil Science at Sam Higginbottom University of Agriculture Technology and Sciences, Allahabad, the area is situated on the south of Allahabad on the right side of the river Yamuna on the South of Rewa Road at a distance of about 6 Km from Allahabad city. It is situated at 25<sup>o</sup>24'23" N latitude, 81<sup>o</sup>50'38" E longitude and at the altitude of 98 meter above the sea level (Table 2).

## **Climate condition in the experimental area**

The area of Allahabad district comes under subtropical belt in the South east of Uttar

Pradesh, which experience extremely hot summer and fairly cold winter. The maximum temperature of the location reaches up to 46<sup>o</sup>C – 48<sup>o</sup>C and seldom falls as low as 4<sup>o</sup>C – 5<sup>o</sup>C. The relative humidity ranged between 20 to 94 percent. The average rainfall in this area is around 1100 mm annually.

The source of nitrogen, phosphorus, neem cake and biofertilizers are as urea, SSP, neem cake, *Rhizobium* culture respectively. The Neem cake applied to some day of before sowing because well decomposed and not direct contact to seed. Basal dose of fertilizer was applied in respective plots according to treatment allocation unifurrows opened by about 5cm. depth before sowing seeds in soil at the same time sowing of seeds was sown on well prepared beds in shallow furrows, at the depth of 5cm, row to row distance was maintained at 30 cm and plant to plant distance was 10 cm, during the course of experiment, observations were recorded as mean values of the data.

## **Results and Discussion**

Table 3 shows the interaction effects of N P K, Neem cake and *Rhizobium* are generally influenced physical and chemical properties of post-harvest soil.

## **Physical properties**

The interaction effects of N P K, Neem cake and *Rhizobium* on bulk density (Mg m<sup>-3</sup>), particle density (Mg m<sup>-3</sup>), % Pore space and Water holding capacity was significant. The maximum bulk density, particle density of after crop harvest soil was recorded as 1.40 Mg m<sup>-3</sup>, 2.51Mg m<sup>-3</sup> respectively in T<sub>0</sub> (I<sub>0</sub>+N<sub>0</sub>+R<sub>0</sub>) @ 0%N P K ha<sup>-1</sup> + @ 0% Neem cake @ ha<sup>-1</sup> + @0% *Rhizobium* ha<sup>-1</sup> and minimum bulk density 1.11 Mg m<sup>-3</sup>, particle density 2.33 Mg m<sup>-1</sup> respectively in @ T<sub>11</sub>- (I<sub>2</sub>+N<sub>1</sub>+R<sub>1</sub>) @100%N P K ha<sup>-1</sup> + @ 100% Neem cake + 100% @ *Rhizobium*. The

highest Pore space and Water holding capacity of soil was recorded 56.62 in @ T<sub>11</sub>-(I<sub>2</sub>+N<sub>1</sub>+R<sub>1</sub>) @100% N P K ha<sup>-1</sup> + @ Neem cake 0.5 t ha<sup>-1</sup> + *Rhizobium* @ 20 g/ Kg seed.

**Chemical properties of post soil**

During the course of study, it was observed that the highest pH was recorded in 7.41 T<sub>0</sub> (I<sub>0</sub>+N<sub>0</sub>+R<sub>0</sub>) @ 0% N P K ha<sup>-1</sup> + @ 0% Neem cake ha<sup>-1</sup> + @ *Rhizobium* ha<sup>-1</sup> and the lowest of 6.97 was recorded with the application of treatment T<sub>11</sub> (I<sub>2</sub>+N<sub>1</sub>+R<sub>1</sub>) @100% N P K ha<sup>-1</sup> + @ 100% Neem cake ha<sup>-1</sup> + @ 100% *Rhizobium*. If we compare the pH of pre sowing soil sample which was 7.44 with that

of after crop harvest soil, there was decrease in pH after crop harvest. Increasing dose of N P K and Neem cake lightly decrease pH of the post-harvest soil. The decrease in pH might be due to higher growth of crops as respiration is more. Respiration evolves carbon dioxide and reacts with water to form carbonic acid in soil.

Organic carbon (%), available nitrogen, phosphorus and potassium (kg ha<sup>-1</sup>) were increased in soil after crop harvests. The chemical properties were significantly affected by different treatment combination of N P K, Neem cake and *Rhizobium*.

**Table.1** Treatment combination of green gram

Treatment	Description	Symbol
T <sub>0</sub>	@ N <sub>0</sub> P <sub>0</sub> K <sub>0</sub> Kg ha <sup>-1</sup> + Neem cake @ 0 t ha <sup>-1</sup> + <i>Rhizobium</i> @ 0 g/ Kg seed	I <sub>0</sub> N <sub>0</sub> R <sub>0</sub>
T <sub>1</sub>	@ N <sub>0</sub> P <sub>0</sub> K <sub>0</sub> Kg ha <sup>-1</sup> + Neem cake @ 0t ha <sup>-1</sup> + <i>Rhizobium</i> @ 20 g/ Kg seed	I <sub>0</sub> N <sub>0</sub> R <sub>1</sub>
T <sub>2</sub>	@ N <sub>0</sub> P <sub>0</sub> K <sub>0</sub> Kg ha <sup>-1</sup> + Neem cake @ 0.5 t ha <sup>-1</sup> + <i>Rhizobium</i> @ 0 g/ Kg seed	I <sub>0</sub> N <sub>1</sub> R <sub>0</sub>
T <sub>3</sub>	@ N <sub>0</sub> P <sub>0</sub> K <sub>0</sub> Kg ha <sup>-1</sup> + Neem cake @ 0.5 t ha <sup>-1</sup> + <i>Rhizobium</i> @ 20 g/ Kg seed	I <sub>0</sub> N <sub>1</sub> R <sub>1</sub>
T <sub>4</sub>	@ N <sub>10</sub> P <sub>20</sub> K <sub>0</sub> Kg ha <sup>-1</sup> + Neem cake @ 0 t ha <sup>-1</sup> + <i>Rhizobium</i> @ 0 g/ Kg seed	I <sub>1</sub> N <sub>0</sub> R <sub>0</sub>
T <sub>5</sub>	@ N <sub>10</sub> P <sub>20</sub> K <sub>0</sub> Kg ha <sup>-1</sup> + Neem cake @ 0 t ha <sup>-1</sup> + <i>Rhizobium</i> @ 20 g/ Kg seed	I <sub>1</sub> N <sub>0</sub> R <sub>1</sub>
T <sub>6</sub>	@ N <sub>10</sub> P <sub>20</sub> K <sub>0</sub> Kg ha <sup>-1</sup> + Neem cake @ 0.5 t ha <sup>-1</sup> + <i>Rhizobium</i> @ 0 g/ Kg seed	I <sub>1</sub> N <sub>1</sub> R <sub>0</sub>
T <sub>7</sub>	@ N <sub>10</sub> P <sub>20</sub> K <sub>0</sub> Kg ha <sup>-1</sup> + Neem cake @ 0.5 t ha <sup>-1</sup> + <i>Rhizobium</i> @ 20 g/ Kg seed	I <sub>1</sub> N <sub>1</sub> R <sub>1</sub>
T <sub>8</sub>	@ N <sub>20</sub> P <sub>40</sub> K <sub>0</sub> Kg ha <sup>-1</sup> + Neem cake @ 0 t ha <sup>-1</sup> + <i>Rhizobium</i> @ 0 g/ Kg seed	I <sub>2</sub> N <sub>0</sub> R <sub>0</sub>
T <sub>9</sub>	@ N <sub>20</sub> P <sub>40</sub> K <sub>0</sub> Kg ha <sup>-1</sup> + Neem cake @ 0 t ha <sup>-1</sup> + <i>Rhizobium</i> @ 20 g/ Kg seed	I <sub>2</sub> N <sub>0</sub> R <sub>1</sub>
T <sub>10</sub>	@ N <sub>20</sub> P <sub>40</sub> K <sub>0</sub> Kg ha <sup>-1</sup> + Neem cake @ 0.5 t ha <sup>-1</sup> + <i>Rhizobium</i> @ 0 g/ Kg seed	I <sub>2</sub> N <sub>1</sub> R <sub>0</sub>
T <sub>11</sub>	@ N <sub>20</sub> P <sub>40</sub> K <sub>0</sub> Kg ha <sup>-1</sup> + Neem cake @ 0.5 t ha <sup>-1</sup> + <i>Rhizobium</i> @ 20 g/ Kg seed	I <sub>2</sub> N <sub>1</sub> R <sub>1</sub>

**Table.2** Mechanical analysis of pre experimental soil

Particulars	Method employed	
Sand (%)	(Bouyoucos,1927)	61.21
Silt (%)		25.36
Clay (%)		13.43
Textural class		Sandy loam
Soil Colour	(Mussel,1908)	
Dry Soil		Pale brown
Wet Soil		Olive brown
Bulk density (Mg m <sup>-3</sup> )	(Muthuaval,1992)	1.32
Particle density (Mg m <sup>-3</sup> )	(Muthuaval, 1992)	2.48
Pore Space (%)	(Black, 1965)	46.42


**Table.3** Chemical analysis of pre experimental soil

Parameters	Method employed	
Soil pH (1:2)	(Jackson,1958)	7.33
Soil EC (dS m <sup>-1</sup> )	(Wilcox, 1950)	0.30
Organic Carbon (%)	(Walleye and Black, 1947)	0.55
Available Nitrogen (Kg ha <sup>-1</sup> )	(Sabbath and Asija, 1956)	287.08
Available Phosphorus (Kg ha <sup>-1</sup> )	(Olsen <i>et al.</i> , 1950)	23.11
Available Potassium (Kg ha <sup>-1</sup> )	(Toth and Prince, 1949)	163.60


**Table.4** Effect of different levels of N P K Neem cake and *Rhizobium* on post-harvest soil properties of Green gram

Treatment combination	Bd (Mg m <sup>-3</sup> )	Pd (Mg m <sup>-3</sup> )	WHC (%)	pH (1:2w/v)	Ec (dSm <sup>-1</sup> )	O.C (%)	N (kg ha <sup>-1</sup> )	P <sub>2</sub> O <sub>5</sub> (kg ha <sup>-1</sup> )	K <sub>2</sub> O (kg ha <sup>-1</sup> )
(I <sub>0</sub> +N <sub>0</sub> +R <sub>0</sub> )	1.40	2.51	48.13	7.41	0.33	0.54	287.13	23.35	168.85
(I <sub>0</sub> +N <sub>0</sub> +R <sub>1</sub> )	1.33	2.47	50.89	7.29	0.32	0.56	290.21	24.53	169.65
(I <sub>0</sub> +N <sub>1</sub> +R <sub>0</sub> )	1.31	2.46	51.33	7.21	0.31	0.59	289.26	25.17	174.47
(I <sub>0</sub> +N <sub>1</sub> +R <sub>1</sub> )	1.23	2.41	53.47	7.13	0.29	0.63	298.50	25.49	175.55
(I <sub>1</sub> +N <sub>0</sub> +R <sub>0</sub> )	1.35	2.47	51.22	7.27	0.32	0.58	295.51	26.19	170.24
(I <sub>1</sub> +N <sub>0</sub> +R <sub>1</sub> )	1.29	2.45	52.85	7.23	0.31	0.60	303.61	27.20	173.51
(I <sub>1</sub> +N <sub>1</sub> +R <sub>0</sub> )	1.25	2.43	54.38	7.17	0.30	0.63	302.71	27.36	176.45
(I <sub>1</sub> +N <sub>1</sub> +R <sub>1</sub> )	1.17	2.38	56.28	7.03	0.29	0.70	307.07	28.25	176.42
(I <sub>2</sub> +N <sub>0</sub> +R <sub>0</sub> )	1.38	2.50	53.25	7.22	0.31	0.61	312.24	30.51	173.74
(I <sub>2</sub> +N <sub>0</sub> +R <sub>1</sub> )	1.19	2.40	56.11	7.08	0.30	0.68	313.87	31.54	175.51
(I <sub>2</sub> +N <sub>1</sub> +R <sub>0</sub> )	1.15	2.36	56.45	7.02	0.29	0.71	314.07	32.48	179.05
(I <sub>2</sub> +N <sub>1</sub> +R <sub>1</sub> )	1.11	2.33	56.62	6.97	0.28	0.74	314.84	33.45	181.05
<b>F- test</b>	S	S	S	NS	NS	S	S	S	NS
<b>S. Me (±)</b>	0.008	0.010	0.04	0.11	0.032	0.012	0.81	0.05	224.37
<b>C. D. at 5%</b>	0.017	0.019	0.07	0.22	0.065	0.024	1.65	0.09	455.93

**Fig.1** Effect of different levels of N P K Neem cake and *Rhizobium* on Bd (Mg m<sup>-3</sup>) and Pd (Mg m<sup>-3</sup>) post-harvest soil properties of Green gram



**Fig.2** Effect of different levels of N P K Neem cake and *Rhizobium* on Post-harvest soil properties of Green gram



The effect of N P K fertilizer on organic carbon (%), available nitrogen, phosphorus, potassium (kg ha<sup>-1</sup>), electric conductivity (dS m<sup>-1</sup>) is significant. The maximum chemical properties of after crop harvest soil was recorded electric conductivity (dS m<sup>-1</sup>), organic carbon (%), available nitrogen, phosphorus, potassium (kg ha<sup>-1</sup>) 0.330, 0.74, 314.84, 33.45, 181. respectively available Potassium (kg ha<sup>-1</sup>), Electric conductivity (ds m<sup>-1</sup>) and pH was found non-significant and Organic carbon (%), Nitrogen (kg ha<sup>-1</sup>), Potassium (kg ha<sup>-1</sup>) and Phosphorus (kg ha<sup>-1</sup>) found to be significant.

It was concluded from trial that the various level of N P K + Neem cake and *Rhizobium* used from in the experiment, the treatment combination T<sub>11</sub>(I<sub>2</sub>+N<sub>2</sub>+R<sub>1</sub>) @100% N P K ha<sup>-1</sup> + @ 100% Neem cake 0.5 t ha<sup>-1</sup> + @ 100% *Rhizobium* 20 g/ Kg seed was found to be the best treatment gave highest benefit of ₹58651 with highest cost benefit ratio 1:2.73 for Green gram it could be recommended for profitable production of Green gram (*Vigna radiata* L.) var. Samrat, it improve soil physical and chemical properties. Integrated nutrient management is better for soil health and Green gram production (Table 4; Figs 1 and 3).

### Acknowledgement

The author are thankful to the Hun' ble Vice Chancellor, HOD, Advisor and non-Teaching staff Department of Soil Science and Agricultural Chemistry, Sam Higginbottom University of Agriculture, Technology And Science Allahabad, U. P., for providing all necessary facilities, clarify studies.

### References

- Ahmed, Z. I., Anjum, M. S. and Raff, C. A. (2006) effect of *Rhizobium* Inoculation on Growth and Nodule Formation of Green gram. *International Journal of Agriculture & Biology*.2:235–237
- Agropedia, (2014-15)
- Anonymous (2011). Agriculture Statistics at a Glance. *Directorate of Economics and Statistics, New Delhi*. The hydrometer as a new method for the mechanical analysis of soils. *Soil Sci*. 23: 393-395.
- Anonymous (2009) Economic survey, Ministry of Finance, Government of India.
- Black, C.A. (1965) Methods of soil analysis. Vol. I. Am. Soc. Agron. Madison, Wisconsin, U.S.A.
- Brady, N.C. and Weil, R.R. (1996) the nature and properties of soils (11th Ed.). Prentice Hall, New York.

- Buoyoucos, G.J. (1952) A recalibration of the hydrometer method for making mechanical analysis of soil, 43, 434.
- Eifediyi, E.K., Mohammed, K.O., Remison, S.U. (2015) effects of neem (*Azadirachta indica* L.) seed cake on the growth and yield of okra (*Abelmoschus esculentus* (L.) Moench) *poljoprivreda*21: (1) 46-52.
- Fisher, R. A. (1950) Technique of Analysis of Variance. Handbook of Agricultural Statistics B-29- 110.
- Hussain, A., Ali, A. and Ijaz, N. R. (2012) Effect of Phosphorus With and Without Rhizobium Inoculation on Nitrogen and Phosphorus Concentration and Uptake by Mungbean (*Vigna radiata* L.). *J. Agric. Res.*, 2012, 50(1).
- (lpr.res.in, 2014-15) ICAR-Indian Institute of Pulses Research, Kanpur-208 024
- Jaiswal, P.C. (2006) Soil, Water and plant Analysis, Manual Practical.
- Jackson, M. L., (1958) Soil chemical analysis, Second edition Indian Reprint, prentice hall of India, New Delhi. Pp498.
- Karpechenko, G. D (1925) Chromosomes of Phaseolinae. *Bull. Appl. Bot.*, 14: 143-148.
- Katyayan A. (2012) Manures, fertilizers and biofertilizers, Fundamentals of agriculture. *Kushal publications and distributors Varanasi*. Vol.-1; p 231-251
- MeenaS.,Swaroop N. and Dawson J. (2016) Effect of integrated nutrient management on growth and yield of green gram (*Vigna radiata* L.) *Agric. Sci. Digest.*, 36 (1) 2016: 63-65
- Murugan, R., Apillai, S.C., Niemsdorff, P.F. and Nanjappan, K. (2011) Effects of combined application of biofertilizers with neem cake on soil fertility, grain yield and protein content of black gram (*Vigna mungo* (L.) Hepper). *World journal of Agricultural sciences* 7 (5): 583-590.
- Muthuaval, P.C., Udayasooriyan, R., Natesan, P.P. and Ramaswami (1992). Introduction to soil Analysis, *Tamilnadu Agriculture University, Coimbatore-641002*.
- Olsen, S. R., Cole, C. V., Watanhe, F. S. and Dean, L. A. (1954) Estimation of available Phosphorus in soils by extraction with sodium bicarbonate U. S. Deptt. Agr. Circ. 939.
- Prasad, K. (2009) a text book field production, ICAR publications, *New Delhi*, pp. 257-258.
- Rajveer, David A. A. and Khadda B. S. (2016) assessment of integrated nutrient on crop growth parameters and yield of green gram (*Vigna radiata* L.). *Society for Scientific Development in Agriculture and Technology*.6277-6279.
- Sahil, S. and P. K. Bandopadhy (2007) Retaining seed vigour and viability of mungbean by drought dressing treatments. *Journal of Food Legumes*. 20: 173-175.
- Subbaiah, B. V. and Asija, C. L. (1956) A rapid procedure for the estimation of available nitrogen in soil. *Current Sciences*; 25:259-260.
- Takase, M., Owusu, L. K. and Sekyere, J. D. (2011) the effects of four sources of irrigation water on soil chemical and physical properties. *Asian Journal of Plant Sciences*; 10: 1, pp 92-96.18.
- Toth, S. J. and Prince, A. L. (1949) Estimation of cation exchange capacity and exchangeable Ca, K and Na content of soil by flame photometer technique. *Soil Sci.*, 67: 439-445.
- Walkley, A. and Black, I. A. (1947) Critical examination of rapid method for determining organic carbon in soils, effect of variance in digestion conditions and of inorganic soil constituents. *Soil sci* pp.632:251.
- Wilcox L.V. (1950) Electrical conductivity, Amer. Water works Assoc. J. 42: pp. 775-776.

#### How to cite this article:

Rajesh and Arun Alfred David. 2017. Response of Neem Cake Rhizobium and Inorganic Fertilizer on Soil Health Growth and Yield of Green gram (*Vigna radiata* L.) var. Samrat. *Int.J.Curr.Microbiol.App.Sci*. 6(7): 815-821. doi: <https://doi.org/10.20546/ijemas.2017.607.100>