

Original Research Article

<https://doi.org/10.20546/ijcmas.2018.706.390>

## Regeneration of Haploids in Oil Seed through Embryonic Microspore

Shahina A. Nagoo<sup>1\*</sup>, GulZaffar<sup>2</sup>, Sabina Naseer<sup>4</sup>, N.A. Dar<sup>3</sup>,  
Sabia Bashir<sup>4</sup>, M. Altaf Wani<sup>4</sup> and Shabina Majid<sup>4</sup>

<sup>1</sup>Division of GPB FoAWadura, SKUAST-K, India

<sup>2</sup>DARS-Budgam SKUAST-K, India

<sup>3</sup>Saffron Research Station, SKUAST-K, India

<sup>4</sup>SKUAST-K, India

\*Corresponding author

### ABSTRACT

Anther Culture or microspore culture is a very important and useful tool in plant breeding for efficient production of haploids and subsequent doubled haploid plants in many species, limited research has been conducted with *Brassica –rapa* as it is more recalcitrant in cell and tissue culture than other Brassicas. In order to access the effect of genotype on microspore embryogenesis in *B –rapa*, two high yielding brown Sarsoon cultivars (KS-101 and KOS-1) were studied for their response to haploid plant regeneration through Anther culture. Flower buds of 2.2 mm to 2.7 mm size representing late uninucleate stage of anther microspores were subjected to low temperature treatment 4°C for 7days and anthers from these buds were cultured on MS medium having 13% sucrose. Anthers were then sub cultured in MS media supplemented with various concentrations of auxins and cytokinins. The best callus development and proliferation was achieved in MS media supplemented with 2,4-D 1mg/l and 0.5 mg/l NAA. Sucrose concentration, cold pre treatment and incubation time influenced embryogenesis. For regeneration of haploid plantlets anther derived calli were transferred to MS full strength medium supplemented with kn92.0mg/l) followed by the use of BA (2.0mg/l) and incubated at 22 ± 20C in light and gave maximum regeneration of 75%. The regeneration shoots (at three leaf stage with length of 5cm) were subculture in rooting media and maximum rooting (30.07%) was achieved when MS medium was supplemented with IBA (0.4mg/l). Root tip mitosis chromosome counts revealed percentage of haploid frequency of KOS-1(41.85) followed by KS-101(38.84%).After 5 days of rooting root samples were taken for confirmation of haploid nature of the plant, Haploid frequency was greater than 65%.

#### Keywords

Haploids, Brassica-  
rapa, Shoot/ Root  
regeneration

#### Article Info

##### Accepted:

22 May 2018

##### Available Online:

10 June 2018

### Introduction

Brown sarson is the major oilseed crop of valley grown on about 50 thousand hacters and fits in rotation with Rice crop. No

significant breakthrough in enhancing the existing productivity levels (7-8 q/ha) and oil quality has been possible due to lack of variability in the available germplasm (Singh *et al* 2007). Operation of self incompatibility

limits production of homozygous inbred lines through conventional breeding procedures, necessitating biotechnological interventions for accelerating breeding progress and generating genetic variability useful for developing high yielding varieties. The efficient production of haploid and double haploid (DH) plants from anther or microspore culture has become an important new tool for *Brassicabreeders*. But limited research has been conducted with Brown Sarson (*Brassicarapa*) as it is more recalcitrant in cell and tissue culture than other *Brassicas*. The present investigation was undertaken to develop a protocol for production of DH Plants in *B.rapa* from androgenic embryos.

### Materials and Methods

Flowered buds 2.5 to 2.7 mm long representing late uninucleate stage were collected from field grown plants of 2 *Brassica compestris* varieties namely KOS-1 and KS-101, subjected to low temperature pretreatment and surface sterilized with 1% sodium hypochloride for 8 to 10 minutes (Gu et al 2004). Anthers were picked from buds and cultured on MS medium supplemented with various concentrations of 2,4-D and NAA for embryogenic callus induction. For regenerations from embryogenic callus MS media supplemented with various concentrations of auxins and cytokinens were

used. Root induction in regenerated shoots was tried using various concentrations of IBA. The cultures were incubated at 22± 2<sup>0</sup> C. Root tip mitotic chromosome counts were used to confirm haploid nature of Anther derived regenerates and double haploidy level induced through use of various concentrations of colchicines was detected cytologically using pollen mother cell (PMC).

### Results and Discussion

Maximum percentage of aseptic culture was achieved in KS-101 followed by KOS-1 genotypes. Callus initiation was maximum (32.62) when anthers were given pretreatment (chilling treatment) temperature of 4<sup>0</sup>C for seven days (Zhang *et al* 2006, Kumari and Singh 2014). Maximum percentage of callus initiation (22.48) was observed in KS-101 genotype followed by KOS-1 (Gu et al 2003) [Table 1].

Maximum number of anthers per 100 buds forming callus was observed in Murashige and Skoog medium 1962, supplemented with 2,4-D (1.0 mg/lit) and NAA (0.5 mg/lit) (36.60). Days taken to initiate callusing was minimum (22.48 days) when induction medium (MS medium) was supplemented with BA (1.5 mg/lit) and 2,4-D (1.0 mg/lit) (Li et al., 2005).

**Table.1** Effect of pre treatment of anthers on callus initiation

Genotype	T1	T2	T3	T4	T5	T6	Mean
KOS-I	23.86 (29.22)	21.88 (27.87)	22.88 (28.55)	30.25 (33.35)	23.02 (28.65)	21.01 (27.26)	<b>23.81</b> <b>(29.15)</b>
KS-IOI	22.03 (27.97)	20.98 (27.24)	21.98 (27.94)	45.78 (42.57)	23.34 (28.88)	19.23 (25.99)	<b>25.55</b> <b>(30.10)</b>

**Table.2** Influence of growth regulators on number of plants regenerated

Growth regulatormg/l	KS-101	KOS-1
BA (0.0)	32.00	48.00
BA(2.0)	69.00	63.00
BA(3.0)	48.00	46.00
Kn(0.0)	44.00	42.00
Kn(2.0)	86.00	72.00
Kn(3.0)	44.00	51.00
Mean	53.83	53.66

**Main effect**      **LSD ≤ (p = 0.01)**      S.E      T<sub>1</sub>=1<sup>0</sup>C  
 Genotype (GT) =      1.44      0.60      T<sub>2</sub> = 2<sup>0</sup>c  
 Temperature (T) =      1.10      0.49      T<sub>3</sub> = 3<sup>0</sup>c  
**Interaction effect**      T<sub>4</sub> = 4<sup>0</sup>c  
 GT x T      =      3.12      1.20      T<sub>4</sub> = 5<sup>0</sup>c      T<sub>6</sub> = 6<sup>0</sup>

**Table.3** Influence of growth regulators on per cent *in vitro* rooting of micro shoots

Growth Regulators	KOS-1	KS-101
IBA(0.0)	24.01(29.33)	18.06(25.14)
IBA(0.2)	25.14(30.09)	20.09(26.62)
IBA(0.4)	26.16(30.75)	23.08(28.71)
IBA(0.6)	21.11(27.35)	19.39(26.12)
IBA(0.8)	19.20(25.98)	18.66(25.58)
Mean	23.12(28.70)	19.89(26.44)

**LSD (p ≤ 0.01)**      S.E ±  
**Main effect**  
 Growth regulator (GR)      =      =      0.78      0.32  
 Genotype (GT)      =      0.88      0.36  
**Interaction effect**  
 GR x GT      =      1.95      0.72

**Table.4** Influence of different treatment combinations and media formulations on haploid frequency (Per cent)

Growth regulators mg l-1	Number of plants regenerated	
	KOS-1	KS-101
BA (0.0)	24.04 (29.36)	18.44 (25.42)
BA (2.0)	30.22 (50.89)	58.25 (49.74)
BS (3.0)	40.21 (39.35)	37.25 (37.61)
Kn (0.0)	30.24 (33.36)	25.24 (30.15)
Kn (2.0)	56.22 (48.57)	54.22 (47.42)
Kn (3.0)	40.22 (39.35)	39.66 (39.03)
Mean	41.85 (40.15)	38.84 (38.23)

**Effect**      **LSD ≤ (p = 0.01)**      S.E ±  
**Main effects**  
 Growth regulator (GR)      =      0.25      0.12  
 Genotype (GT)      =      0.20      0.10  
**Interaction effect**  
 GR x GT      =      0.50      0.24

\* Data in parenthesis are transformed values (Sin<sup>-1</sup> √p)

Highest number of shoots per explants of proliferated cultures (86.00) was achieved (Natalajietal 2006) [Table 2] when MS full strength medium was supplemented with Kn (2.0 mg/lit) followed by the use of BA(2.0 mg/lit)(69.00), highest number of shoots was regenerated in KS-101(53.83)followed by KOS-1(53.66).These shoots devoid of any callus, were sub-cultured on rooting medium containing various concentrations of auxin (IBA). Maximum rooting (30.75%) was achieved when MS medium was supplemented with 0.4mg/lit IBA[Table 3].

Root tip mitosis chromosome counts revealed maximum percentage of haploid frequency in KOS-1(41.85%) followed by KS-101(38.84%) (Natalijaetal 2004)[Table 4]. Diploidization of haploid plants was carried out by using different concentrations of colchicines for different time durations (Dwarkeshetal 2006).Maximum doubling efficiency was obtained when roots of plantlets were submerged in 20mg/lit colchicines for 2 hours and about 40-45 per cent diploidization was achieved. In conclusion, protocol developed for the production of double haploid plants through endrogenesis offers a viable alternative to develop homozygous lines for further use in the development of hybrid/synthetic varieties having higher yield, better quality and tolerance to stresses.

## References

- Dwarkesh, S., Parihar.,Sanjit. A. 2006. Microspore culture for induction of doubled Haploidy in *B. juncea* and *B.rapa* *Journal of Experimental Botany*, 35:1668-1678.
- Gu, H.H., Zhou, W.J., Hagberg, P. 2003. High frequency spontaneous production of doubled haploid plants from microspore culture in *Brassicarapa* ssp. *Chinensis. Ehuphytica*, 134: 239-245.
- Gu, H.H., Tang, G.X., Zhang, G.Q. Zhou. W.J. 2004.Embryogenesis and plant regeneration from isolated microspores of winter cauliflower (*Brassicaoleracea* var. *botrytis*).*Journal of Zhejiang University* (Agriculture Life Science), 30: 34-38.
- Kumari.P and Singh. A.K., 2014.Effect of cold pre-treatment on anther culturein different *Brassicagenotypes*.*AsianJ. Bio. Sci.* 9 (2) Oct.2014, 156-160.
- Li, H.Z., Zhou, W.J. Zhang, Z.J., Gu, H.H., Takeuchi, Y., Yoneyama, K. 2005. Effect of  $\gamma$ -radiation on development yield and quality of micro-tubers *in vitro* in *Solanumtuberosum* L. *BiologiaPlantarium*, 49: 625-628.
- Murashige,T. and Skoog, F. 1962. A revised medium for rapid growth and bioassays with tobacco tissue culture.*Plant physiology*, 15; 473-479.
- Natalija,H., Hegberg,P., Zhou, W.J. 2004. Embryogenesis, callogenesis and plant regeneration from anther culture of spring rape (*B. napus* L.). *PlantGrowth Regulation*, 32; 127-133.
- Singh, V., Singh, S.P., Singh, H., Hegde, D.M. and Tahir, T.A. 2007. Past progress present scenario, nutritional value and strategies to enhance yield potential of rapeseed-mustard: an overview, *Indian Journal of Crop Science* 2(2): 245-257.
- Zhang, G.Q., Zhang, D.Q., Tang, G.X., He, Y., Zhou, W.J.2006. Plant development from microspore-derived embryos in oilseed rapa as affected by chilling, dessication and cotyledon excision. *BiologiaPlantarum*, 50: 180-186.

### How to cite this article:

Shahina A. Nagoo, GulZaffar, Sabina Naseer, N.A. Dar, Sabia Bashir, M. Altaf Wani and Shabina Majid 2018. Regeneration of Haploids in Oil Seed Through Embryonic Microspore. *Int.J.Curr.Microbiol.App.Sci.* 7(06): 3336-3339. doi: <https://doi.org/10.20546/ijcmas.2018.706.390>