

Original Research Article

<https://doi.org/10.20546/ijcmas.2020.912.230>

## Studies on Genetic Variability in Chilli (*Capsicum annum* L.) in the Sundarban Region of West Bengal

Chandan Kumar Mondal<sup>1\*</sup>, Pinaki Acharyya<sup>2</sup> and Pranab Hazra<sup>3</sup>

<sup>1</sup>Ramkrishna Ashram KVK, Nimpith, South 24 Parganas, West Bengal, India

<sup>2</sup>Department of Horticulture, Institute of Agricultural Science, University of Calcutta

<sup>3</sup>Department of Vegetable Science, Faculty of Horticulture, Bidhan Chandra Krishi Viswavidyalaya, Mohanpur, Nadia, West Bengal, India

\*Corresponding author

### ABSTRACT

Chilli is one of the important vegetable as well as spice crop of India. India ranks first in dry chilli production worldwide. Chilli is used throughout the world as a spice and also in the making of beverages and medicines. Since time immemorial chilli is an indispensable part of Indian diet. There are more than 400 cultivars of chilli grown all over the world. Few of them are praised for their red colour because of the pigment 'capsanthin,' while others are known for 'hotness' due to presence of alkaloid 'capsaicin.' It is reported that chilli helps in boosting immune system and lowering of cholesterol. India is rich source of genetic variability of chilli with different quality factors. Limited efforts have been made to release location specific suitable varieties. There is an urgent need and vast scope of evaluating and developing location specific varieties having higher yield potential. In the present investigation thirty seven diverse chilli genotypes were assembled from different parts of India to study their performance in the Sundarban region of West Bengal. The genetic information gathered on yield and yield attributing traits would be of utility in formulation of an efficient breeding programme for the improvement of chilli crop as well as to develop promising genotypes/ hybrids.

#### Keywords

Chilli, Genetic Variability, Sundarban

#### Article Info

##### Accepted:

14 November 2020

##### Available Online:

10 December 2020

### Introduction

Chilli (*Capsicum annum* L.) is one of the most important vegetable as well as spice crop, indispensable in every Indian diet due to its pungency, spicy taste and appealing colour and flavour. It has a predominant position among the spices grown all over India. The production share of chillies among the major spices in India is 25-26 percent (Nandeshwar

and Bharad, 2019). Chilli is praised for its two important qualities, the pungency due to a crystalline acrid volatile alkaloid called Capsaicin and captivating red colour due to the pigment Capsanthin. In India, major chilli producing states are Andhra Pradesh, Orissa, Maharashtra, West Bengal, Karnataka, Rajasthan, Uttar Pradesh and Tamil Nadu (Rajamanickam, 2020). It is reported that majority of the chilli are grown through open

pollinated varieties which have been in cultivation for a long period of time thus leading to low average productivity.

India is an important chilli producing nation in the world sharing 43.4% of global dry chilli production (Anon., 2018a). However, productivity of green chilli is only 11.6 t/ha (produces 35.92 lakh tonnes green chilli from an area of 3.09 lakh hectare land) (Anon., 2018b), which is very low compared to world productivity (14.90 t/ha) (Apeda, 2007). To improve productivity, development of high-yielding varieties is of urgent need. Evaluating and developing location specific varieties having higher yield potential are also important. Limited efforts have been made so far, to release location specific suitable varieties. Selection from the existing variations is the most effective breeding procedure. There are more than 400 cultivars of chilli grown all over the world. India is also a rich source of genetic variability of chilli with different quality factors (Nandi, 1992). To improve yield and other characters, information on genetic variability is of great importance and is a prerequisite for the effective screening of superior accessions (Burton, 1952).

The critical assessment of nature and magnitude of variability in the germplasm stock is the important pre-requisite for formulating effective breeding methods as the genetic improvement of any crop depends on magnitude of genetic variability and the extent of heritability of economically important characters (Jogi *et al.*, 2017). Cramer (2006) suggested that selection based upon multiple characters at the same time may reduce the effectiveness of making improvements in a single trait. It is essential to partition the overall variability into its heritable and non-heritable components with the help of genetic parameters like genotypic co-efficient of variation, heritability and

genetic advance over mean (Vidyashree *et al.*, 2018). Hence the present study was undertaken to determine the extent of genetic variability for important growth and yield characters, interrelationship among the characters and their direct and indirect effects on yield of chilli.

## **Materials and Methods**

**Experimental location:** The field experiment was conducted at the Instructional Farm of Ramkrishna Ashram Krishi Vigyan Kendra, Nimpith, South 24 Parganas, West Bengal consecutively for three years, i.e. during the rabi season of 2008-09, 2009-10 and 2010-11. The Instructional Farm is situated at 88<sup>0</sup>28' East longitude, 22<sup>0</sup>22' North latitude, in the coastal saline zone of West Bengal within the Sundarban ecosystem with an altitude of 3.75m above mean sea level.

**Experimental material:** The experimental material comprised of 37 chilli genotypes assembled from different parts of India (Assam, Nagaland, North Bengal, South Bengal, Pantnagar, Guntur, ICAR-IARI and BCKV-AICRP on Vegetable crops).

**Agronomy:** Forty days old seedlings were transplanted in the main field at a spacing of 60 cm x 45 cm in randomized block design (RBD) with three replications consisting of one row of 30 plants for each entry. The crop was raised as per the recommended package of practices, uniformly applied for each plot.

**Data collection:** Five plants were selected randomly from each plot, tagged and observations were recorded for fifteen growth and yield related traits namely plant height (cm), plant canopy width (cm), number of primary branches per plant, leaf length (cm), leaf width (cm), days to 50 % flowering, fruit length (cm), fruit girth (mm), fruit pedicel length (cm), number of fruits per plant, fresh

fruit weight (g), dry fruit weight (g), number of seeds per fruit, 100 seed weight (mg) and fruit yield per plant (g).

**Statistical analysis:** Average values of each parameter or characters were used for analysis. The mean values were subjected to statistical analysis (ANOVA) as suggested Panse and Sukatme (1967). Phenotypic and genotypic co-efficient of variation, heritability, and genetic advance over mean were calculated following Burton & De-vane (1953), Weber & Moorthy (1952) and Johnson *et al.*, (1955).

### Results and Discussion

Pooled data over three years (2008-09, 2009-10 and 2010-11) have been utilized to study genetic variability among the thirty seven chilli genotypes for the studied fifteen

characters. The analysis of variance, presented in Table 1, clearly suggests significant differences among the genotypes for all the characters (at P=0.01), indicating presence of significant genetic variability in the experimental material. This finding was in accordance with those of Khurana *et al.*, (2003) and Varkey *et al.*, (2005). ‘Year x Genotype’ interaction was significant only for two characters viz. days to 50% flowering and fruit yield per plant, indicating stable expression of most of the characters over the years.

The estimates of mean, range, genotypic co-efficient of variation (GCV), phenotypic co-efficient of variation (PCV), heritability ( $h^2$ ) and genetic advance as percentage of mean (GA) for different characters are presented in Table 2.

**Table.1** ANOVA for growth characters, yield components and yield

Character	Source of Variation				
	Rep.	Year	Genotype	Y x G	Error
DF	2	2	36	72	220
Plant height (cm)	0.827	127.838**	292.112**	5.712	5.759
Plant canopy width (cm)	4.606	3.634	249.095**	1.904	2.909
No. of primary branches /plant	0.478	0.339	12.497**	0.316	0.352
Leaf length (cm)	0.035	0.172	15.293**	0.055	0.106
Leaf width (cm)	0.006	0.028	5.506**	0.011	0.022
Days to 50 % flowering	3.360	6.495	978.284**	5.063**	4.927
Fruit length (cm)	0.036	0.015	13.076**	0.044	0.078
Fruit girth (mm)	0.101	0.442	78.028**	0.268	0.241
Fruit pedicel length (cm)	0.017	0.017	3.329**	0.030	0.046
No. of fruits per plant	2.659	7.073	641.447**	5.653	9.076
Fresh fruit weight (g)	0.005	0.009	2.766**	0.014	0.016
Dry fruit weight (g)	0.002	0.001	0.198**	0.002	0.005
No. of seeds/fruit	0.364	3.214**	3187.201**	0.526	0.580
100 seed weight (mg)	34.201	3.327	11195.133**	9.460	21.922
Fruit yield per plant (g)	7.512	3.681	12035.112**	28.544**	17.736

\* = significant at 5% level

\*\* = significant at 1% level

**Table.2** Genetic variability parameters for different characters

Character	Grand Mean	Range	GCV	PCV	Heritability (h <sup>2</sup> %)	GA as % of mean
Plant height (cm)	50.24	35.89 – 60.24	10.17	11.13	83.49	<b>19.15</b>
Plant canopy width (cm)	44.75	33.47 – 54.30	11.48	11.95	92.32	<b>22.73</b>
No. of primary branches /plant	6.91	5.24 – 9.67	16.57	18.85	77.24	<b>29.99</b>
Leaf length (cm)	7.82	5.48 – 11.54	16.78	17.29	94.08	<b>33.52</b>
Leaf width (cm)	2.87	1.79 – 5.51	27.16	27.68	96.25	<b>54.88</b>
Days to 50 % flowering	54.78	35.67 – 78.00	18.88	19.30	95.66	<b>38.03</b>
Fruit length (cm)	7.85	4.38 – 9.77	15.49	15.86	95.36	<b>31.15</b>
Fruit girth (mm)	10.44	5.64 – 17.11	28.13	28.55	97.13	<b>57.12</b>
Fruit pedicel length (cm)	3.24	2.56 – 5.88	18.56	19.70	88.81	<b>36.04</b>
No. of fruits per plant	46.26	33.78 – 70.90	18.04	19.02	89.96	<b>35.25</b>
Fresh fruit weight (g)	2.79	1.79 – 4.09	19.77	20.24	95.35	<b>39.77</b>
Dry fruit weight (g)	0.75	0.50 – 1.07	19.78	19.88	99.02	<b>40.54</b>
No. of seeds/fruit	66.21	35.76 – 95.99	28.39	28.41	99.84	<b>58.43</b>
100 seed weight (mg)	550.06	492.89 – 611.33	6.40	6.46	98.30	<b>13.08</b>
<b>Fruit yield per plant (g)</b>	<b>125.40</b>	<b>72.35 – 238.35</b>	<b>29.54</b>	<b>29.70</b>	<b>98.93</b>	<b>60.53</b>

It was clear from Table-2 that, the character fruit yield per plant showed maximum range (72.35 – 238.35) followed by number of seeds per fruit (35.76 – 95.99), days to 50 % flowering (35.67 – 78.00), number of fruits per plant (33.78 – 70.90), plant height (35.89 – 60.24) and plant canopy width (33.47 – 54.30). The characters showing wide range of variation provide an ample scope for selecting desired types (Ukkund *et al.*, 2007).

For most of the characters, Phenotypic Coefficient of Variation (PCV) corresponded closely to Genotypic Coefficient of Variation (GCV), which agreed well to the earlier findings of Ukkund *et al.*, (2007). The result suggests that these traits are least affected by environment and selection for these traits on phenotypic performance would be rewarding. In few traits (number of primary branches per plant and fruit pedicel length), the estimates of PCV were greater than GCV. This indicates that the variation for these traits is

not only by genotypes but also due to environment. Hence, selection based on phenotypic performance should not be recommended these traits.

Close correspondence between PCV and GCV estimates was reflected on the broad sense heritability, which were very high (more than 85%) for all the characters except number of primary branches per plant. Burton (1952) suggested that genetic variability along with heritability should be considered for harnessing the maximum and accurate effect of selection. Johanson *et al.*, (1955) opined that heritability in combination with genetic advance would be more reliable for predicting the effect of selection. Genetic advance is the expected genetic progress, possible through selection, over the original population.

Mean genetic advance as percent of mean was high for four traits namely fruit yield per plant (60.53), seeds per fruit (58.43), fruit girth

(57.12) and leaf width (54.88). High value of genetic advance along with moderate GCV and high broad sense heritability was registered for these four traits. To identify the most important characters for selection, it is important to consider the values of GCV, heritability and genetic advance together. Generally high GCV coupled with high broad sense heritability and genetic advance is attributable to additive gene action (Panse, 1957), hence, early generation selection would be helpful for improving these characters. Simple selection will also be effective for the characters like dry fruit weight, fresh fruit weight, days to 50% flowering, fruit pedicel length and fruits per plant which registered moderate GCV, coupled with high heritability and moderate genetic advance.

Moderate to high heritability accompanied with low to moderately low genetic advance was recorded for plant height, plant canopy width, number of primary branches per plant, leaf length, fruit length and 100 seed weight. Low genetic advance coupled with high heritability for the trait plant height was earlier reported by Patil *et al.*, (2008). Improvement of these characters needs selection over several successive years, preferably across locations and over different plantings, because such association of genetic parameters may be attributed to non-additive gene action (Liang and Walter, 1968) and high genotype-environment interaction, hence, simple selection will not be rewarding. Alternately, it suggests heterosis breeding for their amelioration.

## References

- Anonymous (2018a). Tridge: global intelligence of food & agriculture: Chilli Pepper. <https://www.tridge.com/intelligences/other-chili-pepper/production>.
- Anonymous (2018b). Horticultural Statistics at A Glance; Ministry of Agriculture & Farmers' Welfare Department of Agriculture, Cooperation & Farmers' Welfare, Govt. of India. Horticulture Statistics Division; pp 10-11.
- Apeda (2007). <http://apeda.in/agriexchange/Market%20Profile/one/GREEN%20HILLY.aspx>
- Burton G.W. (1952). Quantitative inheritance in grasses. *Proc. 6<sup>th</sup> Int. Grassland Cong.*, 1: 273-283.
- Burton G.W. and De-vane E.H. (1953). Estimating heritability in tall-fescue (*Festuca circundiancaea*) from replicated clonal material. *Agron. J.* 45: 478-481.
- Cramer C.S. (2006). Onion trait heritability and response from selection. *J. Amer. Soc. Hort. Sci.*, 131(5): 646-650.
- Jogi, M.Y., Madalageri, M.B., Mallimar, M., Bawoor, S., Mangi, V. and Porika, H., (2017). Genetic Variability Studies in Chilli (*Capsicum annum L.*) for Growth and Early Yield, *Int. J. Pure App. Biosci.* 5(5): 858-862.
- Johanson H.W., Robinson, H.F. and Comstock, R.E. (1955). Estimates of genetic and environmental variability in Soyabean. *Agron. J.*, 47(7): 314-315.
- Khurana, D.S., Singh, P. and Hundal, J. S. (2003). Studies on genetic diversity for growth, yield and quality traits in chilli (*C. annum L.*). *Indian J. Hort.*, 60(3): 277-282.
- Liang G.H.L. and Walter, T.H. (1968). Heritability estimates and gene effects for agronomic traits in grain sorghum (*Sorghum vulgare*). *Crop Sci.*, 8: 77-80.
- Nandeshwar V.N. and Bharad S.G. (2019). Effect of planting geometry and fertigation levels on growth, yield and quality of chilli. *J Krishi Vigyan*, 8 (1): 63-69.
- Nandi A. (1992). Genetic variability in chilli

- (*Capsicum annuum* L.). *Indian journal of Coca Arecanut and Spices.*, 16 (3/4): 140-145.
- Panse V. G. and Sukhatme, P. V. (1967). Statistical methods for agricultural workers, Indian Council of Agricultural Sciences, New Delhi.
- Patil, S.D., Bidari, B.I., Shashidhara, G.B. and Hegde, N.K. (2008). Genetic variability in chilli (*Capsicum annuum* L.) genotypes. *Asian J. Hort.*, 3(2): 310-312.
- Rajamanickam C. (2020). Assessment of Chilli (*Capsicum annuum* L.) hybrids for growth and yield characters. *J Krishi Vigyan*, 8 (2): 12-15.
- Ukkund K.C., Madalageri, M.B., Patil, M.P., Mulage, R. and Kotikal, Y.K. (2007). Variability studies in green chilli (*Capsicum annuum* L.). *Karnataka J. Agric. Sci.*, 20(1): 102-104.
- Varkey J., Saiyed, M.P., Patel, J.S. and Patel, D.B. (2005). Genetic variability and heritability in chilli. *J. Maharashtra Agric. Univ.*, 30(3): 346-347.
- Vidyashree A.B., Allolli, T.B., Ravi, Y. and Mulge, R. (2018). Genetic Variability Studies in Byadagi Chilli (*Capsicum annuum* L.), *Int. J. Pure App. Biosci.* 6(5): 1120-1125.
- Weber C. R. and Moorthy H. R. (1952). Heritable and non-heritable relationship and variability of oil content and agronomic characters in the F2 generation of soybean crosses. *Agron. J.*, 44: 202-209.

**How to cite this article:**

Chandan Kumar Mondal, Pinaki Acharyya and Pranab Hazra. 2020. Studies on Genetic Variability in Chilli (*Capsicum annuum* L.) in the Sundarban Region of West Bengal. *Int.J.Curr.Microbiol.App.Sci.* 9(12): 1938-1943. doi: <https://doi.org/10.20546/ijcmas.2020.912.230>