

Original Research Article

<https://doi.org/10.20546/ijcmas.2020.912.179>

Comparative Study of Energy Use Pattern of Conventional Farming and Organic Farming in Rice Production in Visakhapatnam District of Andhra Pradesh State

S. Shrine* and K. Umadevi

Department of Agricultural Economics, Agricultural College, Acharya N. G. Ranga
Agricultural University, Bapatla, India

**Corresponding author*

ABSTRACT

Keywords

Conventional Farming,
Organic Farming

Article Info

Accepted:
12 November 2020
Available Online:
10 December 2020

The study examined the energy use pattern of conventional farming and organic farming in rice production in Visakhapatnam district. Fertilizer was found to be the dominant source of energy in both farming methods of rice production. The total energy utilized for rice production was highest in conventional farming compared to organic farming. The energy use efficiency was highest in organic farming than conventional farming.

Introduction

The important sector in Indian economy is the agriculture and it is one of the most energy consuming economies. During 2017-18, rice production in India was 111.01 million tonnes and in India, Andhra Pradesh ranks 5th in rice production. In Andhra Pradesh, rice is cultivated in 22.18 lakh hectares and production and productivity were about 126.91 lakh tonnes and 5722 kg ha⁻¹ (2017-18) respectively. In Andhra Pradesh, Visakhapatnam district has a production of 3.43 lakh tonnes in 109 thousand hectares. In India, the rice is cultivated in different farming methods such as conventional

farming, organic farming, zero budget natural farming etc., The energy use pattern for rice crop has varied under different farming methods. The use of energy in crop production depends on the availability of energy sources and capacity of farmers.

Conventional farming refers to farming systems which include the use of synthetic chemical fertilizers, pesticides, herbicides and other inputs, genetically modified organisms, concentrated animal feeding operations, heavy irrigation and intensive tillage. Organic farming involves cultivation of crop using biological materials, avoiding synthetic substances to maintain soil fertility and

ecological balance. The energy consumption varies from each other in both farming methods. It is invested in various forms such as mechanical power, human labour, animal draft, fertilizer, pesticides etc. Ploughing consumed highest amount of energy which accounted to 20.58 per cent of the total energy utilized for all operations in paddy cultivation (Kumar and Hugar, 2011). Sufficient availability of the right energy and its effective and efficient use are prerequisites for improved agricultural production. It was realized that crop yields and food supplies are directly linked to energy (Stout, 1990).

Energy is invested in various forms such as mechanical power, human labour, animal draft, chemical fertilizer, pesticides, herbicides etc., So, the present study is based on energy consumption in two different farming methods in rice production.

Materials and Methods

The present study was carried out in Visakhapatnam district of Andhra Pradesh (AP) state during 2017-18. A multi-stage random sampling method was adopted for the selection of mandals, villages and farmers. In first stage, four mandals namely, Anakapalle, Chodavaram, Paderu, Hukumpeta were selected in Visakhapatnam district, as the farmers were practicing conventional farming in Anakapalle and Chodavaram mandals and Organic farming in Paderu and Hukumpeta mandals. In second stage, two villages from each mandal were selected based on the maximum number of farmers practicing three farming methods. In third stage, a total sample size of 109 respondents out of 625 was selected based on Cochran's formula for the present study. The respondents following conventional farming and organic farming were selected based on the proportional total.

$$n_0 = \frac{Z^2 pq}{e^2}$$

$$n = \frac{n_0}{1 + \frac{n_0 - 1}{N}}$$

Where, n= sample population

N= Total population

Z = Critical value

e = sampling error

p= 0.5, q= 1- 0.5

Inputs and output energy was calculated by converting physical units of inputs and outputs into respective energy units by using appropriate energy equivalents presented in Table 2 to find out the energy use pattern.

Agriculture is not only a consumer of energy but also a producer of energy in the form of output. The efficiency of energy can be calculated by ratio of output energy to input energy.

Energy productivity= Rice output (Kg ha⁻¹)/ Energy input (MJ ha⁻¹)

Specific energy= Energy input (MJ ha⁻¹)/ Rice output (Kg ha⁻¹)

Net energy= Energy output (MJ ha⁻¹)- Energy input (MJ ha⁻¹)

Energy intensiveness= Energy input (MJ ha⁻¹)/ Cost of cultivation (Rs ha⁻¹)

Energy use efficiency= Energy output (MJ ha⁻¹)/ Energy input (MJ ha⁻¹)

Results and Discussion

The total operation-wise energy consumption in organic farming was 13978.84 MJ/ha. Among all the inputs, organic fertilizers and manures (8823.42 MJ/ha.) consumed highest energy followed by human (1332.90 MJ/ha.), machinery (1322.12 MJ/ha.), seed (1089.00 MJ/ha.), biocides (940.06 MJ/ha.) and animal (471.34 MJ/ha.) (Table 1-4).

Table.1 Sampling frame of respondents

S. No.	Mandals	Villages	Population size	Per cent to total population	Sample size	Per cent to total sample
1.	Anakapalle	Dibbapalem	102	13.04	18	13.04
		Venkupalem	98	12.53	17	12.53
2.	Chodavarm	Narsapuram	104	13.29	18	13.29
		Lakshmipum	92	11.76	16	11.76
3.	Paderu	Thumpada	22	2.81	4	2.81
		Vanajangi	57	7.28	10	7.28
4.	Hukumpeta	Rangasheela	92	11.76	16	11.76
		Bodigaputtu	58	7.41	10	7.41
	TOTAL		625		109	

Table.2 Energy equivalents of different inputs and outputs

S. NO.	ENERGY SOURCE	UNITS	ENERGY EQUIVALENTS (MJ)
A.	Inputs		
1.	Human labour		
	A. Man	h	1.96
	B. Woman	h	1.57
2.	Animal labour		
	A. Bullocks	Pair- hour	10.10
3.	Machine labour	H	62.70
4.	Fertilizers		
	A. Nitrogen	kg	66.44
	B. Phosphorous	kg	12.44
	C. Potash	kg	11.15
	D. Sulphur	kg	1.20
7.	Organic fertilizers		
	Farmyard manure	kg	0.33
	Commercial compost	kg	10.50
	Green manure	kg (seed)	15.90
	Bio- fertilizers	L	2.98
8.	Chemicals		
	A. Herbicides	kg	238.00
	B. Pesticides	kg	199.00
	C. Fungicide	kg	92.00
9.	Bio- fungicide	L	10.10
	Bio- insecticides	L	21.60
10.	Seeds	Kg	14.70
B.	Output		
	Rice	Kg	17.00

L= Litres, kg= Kilogram, h= hour

Source: Sefeedpariet *al.* (2012), Bilaliset *al.* (2013)

Table.3 Energy use pattern of rice production (MJ ha⁻¹)

Particulars	Conventional farming (N= 69)		Organic Farming (N= 40)	
	Energy (MJ/ha.)	Per centage	Energy (MJ/ha.)	Per centage
Inputs				
Human	1778.42	7.71	1332.9	9.53
Machine	4552.46	19.74	1322.12	9.45
Animal	-	-	471.34	3.37
Seed	1046.37	4.53	1089.00	7.79
Natural fertilizers and manures	-	-	-	-
Chemical fertilizers and manures	15351.87	66.58	-	-
organic fertilizers and manures	-	-	8823.42	63.11
Plant protection chemicals	326.48	1.41	-	-
Biocides	-	-	940.06	6.72
Total	23055.6	100	13978.84	100.00
Output				
Yield	64771.87		75796.87	

Table.4 Energy indices for rice production

Parameter	Conventional farming	Organic farming
Energy productivity (kg MJ⁻¹)	0.17	0.38
Specific energy (MJ kg⁻¹)	5.57	2.63
Net energy (MJ ha⁻¹)	47468.63	60683.019
Energy use efficiency	3.05	5.27

In two different farming methods for rice production, conventional farming consumed maximum amount of energy (23055.6 MJ/ha.) compared to organic farming (13978.84 MJ/ha.). Among all the operations in two different farming methods, fertilizer and manures consumed maximum amount of energy. These findings are in line with the findings of Pradhan *et al.*, (2015) who reported that fertilizer consumed highest amount of energy in rice production. The

maximum yield energy estimated for rice production was in organic farming (75796.87 MJ/ha.) compare to conventional farming (64771.87 MJ/ha.).

From Table 4, in conventional farming, the energy productivity was found to be 0.17 kg/MJ. That means 0.17 kg grain output was obtained per unit energy. The specific energy was found to be 5.57 MJ/kg. Initiating that, to produce one kg of grain output 5.57 MJ

energy was consumed. The net energy was found to be 47468.63 MJ ha⁻¹. The energy use efficiency was found to be 3.05 which indicates the efficiency of rice production.

In organic farming, the energy productivity was found to be 0.38 kg/MJ. That means 0.38kg grain output was obtained per unit energy. The specific energy was found to be 2.63 MJ/kg. That means, to produce one kg of grain output 2.63 MJ energy was consumed. The net energy was found to be 60683.019 MJ ha⁻¹. The energy use efficiency was found to be 5.27 which indicate the efficiency of rice production.

The energy use efficiency was found to be higher in organic farming compared to conventional farming because in organic farming, less input energy was consumed per unit output energy. The net energy was found to be high in organic farming than conventional farming.

In conclusion, the total energy consumption was highest in conventional farming compared to organic farming. The difference in energy consumption is mainly due to (chemical/organic) fertilizers and manures in the two farming methods. The energy use efficiency is highest in organic farming compared to conventional farming. The net energy was found to be high in organic farming compared to conventional farming.

References

- Kumar, P.P.S and Hugar, L.B. 2011. Economic analysis of energy use in paddy cultivation under irrigated situations. *Karnataka Journal of Agricultural Sciences*. 24(4):467-470.
- Stout, B. A. 1990. Handbook of Energy for World Agriculture. *Elsevier Applied Science*, London.
- Sefeepari, P., Rafiee, S., Komleh, S. H. P and Ghahderijani, Md. 2013. A sourcewise and operationwise energy use analysis for corn silage production, a case study of Tehran province, Iran. *International Journal of Sustainable Built Environment*. 1: 158- 166.
- Bilalis, D., Kamariari, P. E., Karkanis, A., Efthimiadou, A., Zorpas, A and Kakabouki, L. 2013. Energy inputs, output and productivity in organic and conventional maize and tomato production, under Mediterranean conditions. *NotulaeBotanicaeHortiAgrobotanici Cluj-Napoca*. 41 (1): 190- 194.
- Pradhan, P., Naik, R.K., Manisha sahu and Cholesh, T. 2015. A study on the energy use pattern and cost of production under transplanted paddy production system in Chhattisgarh, India. *International Journal of Engineering Research & Technology*. 4 (7): 1014-1018.

How to cite this article:

Shrine, S. and Umadevi, K. 2020. Comparative Study of Energy Use Pattern of Conventional Farming and Organic Farming in Rice Production in Visakhapatnam District of Andhra Pradesh State. *Int.J.Curr.Microbiol.App.Sci*. 9(12): 1505-1509.
doi: <https://doi.org/10.20546/ijcmas.2020.912.179>