

Original Research Article

<https://doi.org/10.20546/ijcmas.2020.909.369>

Genetic Variability Studies for Yield and Quality Traits in Chickpea (*Cicer arietinum* L.)

T. Anusha^{1*}, A. Trivikrama Reddy¹, V. Jayalakshmi² and S. Khayum Ahammed²

¹Department of Genetics and Plant Breeding, Agricultural College, Mahanandi-518502, India
²Regional Agricultural Research Station, Nandyal-518501, India

*Corresponding author

ABSTRACT

Keywords

Chickpea, Genetic variability, Phenotypic coefficient of variation, Genotypic coefficient of variation, Heritability, Genetic advance as per cent of mean

Article Info

Accepted:
20 August 2020
Available Online:
10 September 2020

The present investigation was carried out with 25 chickpea genotypes to study the genetic variability parameters for yield and quality traits of chickpea and observed highly significant differences among the genotypes for all the eleven traits studied viz., plant height, number of branches plant⁻¹, days to 50% flowering, days to maturity, number of pods plant⁻¹, 100 seed weight, harvest index, protein content, iron content, zinc content and seed yield plant⁻¹. Genetic variability parameters showed moderate to high estimates of phenotypic coefficient of variation and genotypic coefficient of variation, with high heritability in broad sense coupled with high genetic advance as per cent of mean for the traits viz., plant height, number of pods plant⁻¹, 100 seed weight, seed yield plant⁻¹ and iron content and thus, indicating that these characters exhibited wide range of variability. Heritability in these characters was due to additive gene effects and selection may be effective. This will help the breeders for direct selection of plants on the basis of phenotypic expression.

Introduction

Chickpea (*Cicer arietinum* L.) is highly self-pollinated diploid annual leguminous plant. It is the third most important pulse crop in the world after faba bean and field pea (Singh *et al.*, 2018). India is the world's largest producer of chickpea with annual production of 10.13 million tonnes from an area of 9.44 million ha with productivity of 1073 kg ha⁻¹ (Directorate of Economics and Statistics,

2019). Other important countries growing chickpea are Pakistan, Mexico, Turkey, Canada, Iran, Australia, Tanzania, Ethiopia, Spain and Burma. In Andhra Pradesh, chickpea is grown in an area of 4.78 lakh ha producing 2.43 lakh tonnes with productivity 506 kg/ha (Directorate of Economics and Statistics, 2019).

Chickpea is highly nutritious crop and is a protein- rich supplement to cereal based diets.

Chickpea seeds contain on an average 23% protein, 64% total carbohydrates (47% starch, 6% soluble sugar), 5% fat, 6% crude fibre and 2% ash and also contain micro nutrients like phosphorus (340 mg/100 g), calcium (160 mg/100 g), magnesium (140 mg/100 g), iron (5 mg/100 g), zinc (4.1 mg/100 g) (Jukanti *et al.*, 2012). Since, cereals combined with pulses are a major diet for billions of people around the world, chickpeas are being targeted in many countries so that malnutrition problem can be controlled to some extent.

Genetic improvement of any crop mainly depends on the magnitude of genetic variability present in the breeding material and to extent to which the yield and quality characters are heritable from generation to generation. The estimates of genotypic and phenotypic coefficient are necessary for understanding the influence of environment on different traits.

Heritability and genetic advance as *per cent* of mean is a reliable tool in the selection programme to get a clear picture of the scope of improvement of various characters through selection. The identification of chickpea genotypes rich with protein and micronutrients along with good yield help the breeders to identify donars for the future breeding programme. Keeping this in view, the present investigation was carried out with the following objective to estimate the extent of genetic variability for yield and quality traits in chickpea.

Materials and Methods

The present investigation was carried out with 25 genotypes comprising of 20 advanced *desi* breeding lines and 5 check lines at Regional Agricultural Research Station, Nandyal during *rabi* 2019-20 in a Randomized Block Design (RBD) with three replications and

spacing of 30 cm x 10 cm was adopted. Recommended agronomical and plant protection practices for chickpea crop were followed. Observations were recorded for 11 traits *viz.*, plant height, number of branches plant⁻¹, number of pods plant⁻¹, days to 50% flowering, days to maturity, 100 seed weight, harvest index, protein content, iron content, zinc content and seed yield plant⁻¹ as per standard procedures.

For the estimation of protein content, mature seeds from each genotype in each replication were powdered and 0.2 g of powdered seed sample was digested with concentrated H₂SO₄ for determination of nitrogen content in per cent by using Micro – Kjeldhal digestion and distillation method as given by AOAC (1965).

The crude protein content in seeds was obtained by multiplying the determined nitrogen content with 6.25. A known quantity of 0.2 g powdered sample was digested with triacid mixture (nitric acid, sulfuric acid and perchloric acid @ 9:4:1) to determine the iron content and zinc content by using Atomic Absorption Spectrophotometer.

Analysis of variance was carried out for each character separately as per the method outlined by Panse and Sukhatme (1985). The genotypic coefficient of variation (GCV) and phenotypic coefficient of variation (PCV) were estimated as per the formula suggested by Burton (1952) while heritability in broad sense by Lush (1940) and genetic advance as per cent of mean by Johnson *et al.*, (1955 a).

Results and Discussion

Analysis of variance revealed highly significant differences among the genotypes for all the 11 characters studied (Table 1) and shown a wide range of variability for the characters considered.

Range of variability

The range of variability for plant height (29.0 cm to 47.5 cm), number of branches plant⁻¹ (11.2 to 18.4), days to 50 % flowering (35.0 days to 53.3 days), days to maturity (81.0 days to 90.7 days), number of pods plant⁻¹ (23.3 to 45.2), 100 seed weight (20.9 g to 36.0 g), harvest index (50.5% to 64.8%), protein content (18.1% to 22.1%), iron content (6.5 mg/100g to 14.4 mg/100g), zinc content (3.0 mg/100g to 5.3 mg/100g) and seed yield plant⁻¹ (4.8 g to 12.0 g) were observed in Table 2.

The variation among the chickpea genotypes in this study would provide ample opportunities for the genetic improvement of the crop through direct selection of genotypes or through hybridization using as parents possessing the desirable traits.

Phenotypic and Genotypic coefficient of variation

The estimates of phenotypic coefficient of variation (PCV) for all 11 characters were observed to be higher than the estimates of genotypic coefficient of variation (GCV) (Table 2), indicating the influence of environment on the expression of these characters which was also reported by Raju *et al.*, (2017), Srivastava *et al.*, (2017) and Mohammed *et al.*, (2019). PCV and GCV values of more than 20% are considered to be high, values between 10 – 20% as medium and values less than 10% as low (Sivasubramanian and Menon, 1973).

In this study, iron content recorded high values of PCV and GCV, indicating the presence of wide variability among the genotypes studied for this trait. On the other hand, days to 50% flowering, days to maturity, harvest index and protein content showed low levels of PCV and GCV,

signifying limited scope for further genetic improvement through selection. The traits *viz.*, plant height, number of branches plant⁻¹, number of pods plant⁻¹, 100 seed weight, zinc content showed moderate magnitude of PCV and GCV. Seed yield plant⁻¹ exhibited high PCV and moderate GCV values.

These findings were in conformity with findings of Jayalakshmi *et al.*, (2018) for iron content, Jayalakshmi and Trivikrama Reddy (2018) for protein content and zinc content, Raju *et al.*, (2017) for days to 50 % flowering, plant height and number of branches plant⁻¹ per plant. Similarly, Banik *et al.*, (2018), Mohammed *et al.*, (2019) and Mohan and Thiyagarajan (2019) for days to maturity, Kumar *et al.*, (2018) for number of pods plant⁻¹

Heritability in broad sense and genetic advance as per cent of mean

High heritability in broad sense estimates was observed in Table 2 for characters *viz.*, number of pods plant⁻¹ (93.6%), iron content (93.0%), days to 50% flowering (91.7%), 100 seed weight (91.6%), plant height (90.6%), days to maturity (90.1%), and seed yield plant⁻¹ (73.0%) and thus indicated the presence of a considerable proportion of total genetic variance which includes fixable and non-fixable variances.

Moderate heritability estimates were recorded for harvest index (57.3%), zinc content (57.1%), number of branches plant⁻¹ (49.0%) and protein content (39.7%). Similar findings were also reported by Desai *et al.*, (2015), Raju *et al.*, (2017), Banik *et al.*, (2018) and Singh *et al.*, (2018) for plant height, days to 50 % flowering, number of pods plant⁻¹, 100 seed weight and seed yield plant⁻¹; Mohan and Thiyagarajan (2019) for number of branches plant⁻¹ and protein content; Srivastava *et al.*, (2017) for harvest index.

Table.1 Analysis of variance for 11 characters in 25 genotypes of chickpea

S.No	Characters	Mean sum of squares		
		Replications	Treatments	Error
		(df : 2)	(df: 24)	(df: 48)
1	Plant height (cm)	10.1	58.1**	1.9
2	Number of branches plant ⁻¹	8.6	8.7**	2.2
3	Days to 50% flowering	46.3	47.0**	1.4
4	Days to maturity	40.4	17.3**	0.6
5	Number of pods plant ⁻¹	35.6	131.6**	2.9
6	100 seed weight (g)	1.1	40.8**	1.2
7	Harvest index (%)	0.3	37.3**	7.4
8	Protein content (%)	8.4	3.4**	1.1
9	Fe content (mg/100g)	1.6	16.8**	0.4
10	Zn content (mg/100g)	1.2	1.0**	0.2
11	Seed yield plant⁻¹ (g)	2.2	9.0**	1.0

** Significant at 1% level

Table.2 Estimates of genetic parameters for 11 characters in 25 chickpea genotypes

S.No.	Characters	General Mean	Range		GCV (%)	PCV (%)	Heritability (Broad sense) (%)	Genetic Advance as per cent of mean
			lowest	Highest				
1	Plant height (cm)	36.5	29.0	47.5	11.8	12.4	90.6	23.2
2	Number of branches plant ⁻¹	14.5	11.2	18.4	10.1	14.4	49.0	14.6
3	Days to 50% flowering	46.3	35.0	53.3	8.4	8.8	91.7	16.6
4	Days to maturity	84.6	81.0	90.7	2.8	2.9	90.1	5.5
5	Number of pods plant ⁻¹	33.9	23.2	45.2	19.3	20.0	93.6	38.5
6	100 seed weight (g)	26.5	20.9	36.0	13.7	14.3	91.6	27.0
7	Harvest index (%)	58.0	50.5	64.8	5.4	7.2	57.3	8.5
8	Protein content (%)	20.1	18.1	22.1	4.3	6.8	39.7	5.6
9	Fe content (mg/100g)	10.6	6.5	14.4	21.9	22.7	93.0	43.5
10	Zn content (mg/100g)	3.0	2.6	5.3	12.4	16.5	57.1	19.3
11	Seed yield plant⁻¹ (g)	8.9	4.8	12.0	18.4	21.5	73.0	32.4

GCV: Genotypic coefficient of variation; PCV: Phenotypic coefficient of variation

Genetic advance as per cent of mean was high for iron content (43.5%), number of pods plant⁻¹ (38.5%), seed yield plant⁻¹ (32.4%), 100 seed weight (27.0 %) and plant height (23.2%) thus showed that these characters were governed by additive genes and selection will be rewarding for improvement of these characters. Moderate genetic advance as *per cent* of mean was observed for zinc content (19.3%), number of branches plant⁻¹ (14.6%) and days to 50% flowering (16.6%).

Low genetic advance as *per cent* of mean was observed for harvest index (8.5%), protein content (5.6%) and days to maturity (5.5%) and thus indicated that these characters were governed by non-additive genes. Similar findings were also reported by Raju *et al.*, (2017) and Singh *et al.*, (2018) for 100 seed weight, plant height, days to 50% flowering; Jayalakshmi *et al.*, (2018) for iron content; Desai *et al.*, (2015) for protein content.

In the present investigation, plant height, number of pods plant⁻¹, 100 seed weight, iron content and seed yield plant⁻¹ showed high estimates of broad sense heritability coupled with high genetic advance as per cent of mean indicating that these characters could be prominently governed by additive gene action and selection of these traits may be more effective for desired genetic improvement.

It was supported by similar findings of Desai *et al.*, (2015), Singh *et al.*, (2018), Banik *et al.*, (2018) for 100 seed weight, number of pods plant⁻¹ and plant height; Jayalakshmi *et al.*, (2018) for iron content.

In conclusion the present studies revealed that iron content showed high PCV and GCV whereas, plant height, number of branches plant⁻¹, number of pods plant⁻¹, 100 seed weight, zinc content showed moderate magnitude of PCV and GCV. Seed yield plant⁻¹ exhibited high PCV and moderate

GCV values. The traits *viz.*, plant height, number of pods plant⁻¹, 100 seed weight, iron content and seed yield plant⁻¹ showed high estimates of broad sense heritability coupled with high genetic advance as per cent of mean

The assessment of genetic parameters like phenotypic and genotypic coefficient of variation, heritability and genetic advance as per cent of mean indicated that selection must be done in the characters like plant height, number of pods plant⁻¹, 100 seed weight, seed yield plant⁻¹ and iron content for improving the yield and nutritional traits to generate genetic variability followed by selection in further generations to identify superior segregants for these characters.

References

- AOAC. 1965. Official methods of analysis of the Association of Official Agricultural Chemists. 8th edition. Minnosta.
- Banik, M., Deore, G.N., Mandal, A.K and Mhase, L.B. 2018. Genetic variability and heritability studies in chickpea (*Cicer arietinum* L.). *Current Journal of Applied Sciences and Technology*. 31(1): 1-6.
- Burton, G.W. 1952. Quantitative inheritance in grasses. *Proceedings of 6th Grassland Congress Journal*. 1: 277-281.
- Desai, K., Tank, C.J., Gami, R.A., Patel, A.M and Patel, C. 2015. Character association analysis for yield, its components and nutritional quality in chickpea (*Cicer arietinum* L.). *Research Journal of Agricultural Sciences*. 6(6): 1280-1283.
- Directorate of Economics and Statistics, 2019. Ministry of Agriculture Cooperation and Farmers Welfare (GOI). *Agricultural statistics at a glance 2019*. pp. 62-63.

How to cite this article:

Anusha, T., A. Trivikrama Reddy, V. Jayalakshmi and Khayum Ahammed, S. 2020. Genetic Variability Studies for Yield and Quality Traits in Chickpea (*Cicer arietinum* L.). *Int.J.Curr.Microbiol.App.Sci.* 9(09): 2995-3000. doi: <https://doi.org/10.20546/ijcmas.2020.909.369>