

Original Research Article

Biofertilizers for Growth and Establishment of Alphonso Mango Grafts under Nursery Condition

T. H. Shankarappa^{1*}, R. Narayana Reddy², B. Subramanyam³, A. Sreenatha³ and N. Aswathanarayana Reddy³

¹Regional Horticultural Research and Extension Centre, GKVK, Bengaluru, 560 065, India

²Staff Training Unit, UAS, Hebbal Campus, Bengaluru, 560 024, India

³Horticultural Research & Extension Centre, Hogalagere, Kolar 563 138, India

*Corresponding author

ABSTRACT

Experiment on establishment of mango grafts (Root stock: Totapuri (Bangalora) and Scion: Alphonso) was conducted in green house at Horticulture Research station, Hogalagere, Kolar during the period 2015-16. The grafts were planted in potting mixture; soil with farm yard manure and sand in the ratio of 2:1:1 (Soil: FYM; Sand) taken in polythene bags of size 12 cm x 18 cm. Applied with biofertilizers such as *Azospirillum* spp., Phosphorus solubilizing bacteria (PSB), *Pseudomonas fluorescens*, @ 10 g per graft (single / dual / combinations 10g only) plus Vesicular Arbuscular Mycorrhiza (VAM), soil based biofertilizer @10 g. The moisture content of soil in poly bags was maintained at field capacity. A total of fifteen treatments were tested in three replications. The potting mixture had pH - 6.95, Organic Carbon - 0.86 %, Available N- 228 kg ha⁻¹, Available P- 35 kg ha⁻¹ and available Potassium 310 kg ha⁻¹. The results at the end of 150 days showed that the inoculation of consortium of three biofertilizers; *Azospirillum*, *Pseudomonas fluorescens* and VAM had produced highest plant height (68.23 cm), stem girth (3.83 cm) and number of leaves (18.33 no.). The inoculation of consortium of three biofertilizers was found to be on par with inoculation of consortium containing four biofertilizers but found to be superior over dual inoculation, single inoculation and over control. The inoculation of consortia of these three biofertilizers showed better nutrient uptake and growth at the end of 150 days in nursery.

Keywords

Mango grafts, alphonso, biofertilizer, growth, nutrient uptake

Introduction

Mango (*Mangifera indica* L.) belonging to family anacardiaceae is one of the most important fruit of India. Mango occupies 22% of the total area under fruits comprising of 1.2 million hectares, with a total production of 11 million tonnes. Uttar Pradesh and Andhra Pradesh, Telangana are having the largest area under mango each with around 25% of the total area followed by Bihar, Karnataka, Kerala and Tamil Nadu

(Mehta, 2017). In 2014-15, Karnataka state had produced 1755.56 thousand MT of mango with a productivity of 9.39 MT/ha (National mango database, 2015). The expansion of area under cultivation of this crop is increasing annually in Karnataka because mango is well adapted to tropical and sub-tropical climate of Karnataka which is supported by loamy, alluvial, well drained, aerated and deep soils rich in

organic matter with a pH range of 5.5 to 7.5. Although mango can be propagated from seed, vegetative propagation is practiced commonly. Plants are generally propagated by grafting onto seedling rootstocks. Grafted trees usually produce fruit in 3 to 5 years in dry areas, while seedling trees usually take at least five years to come into bearing.

One of the limitations in mango cultivation is the production of good quality grafted plants and their establishment in nursery. It is very much required to ensure survivability and further establishment in main fields. These limitations can be narrowed by use of biofertilizers such as *Azospirillum*, phosphorus solubilizing bacteria, *Pseudomonas fluorescens* and Mycorrhizae. Biofertilizers such as *Azospirillum*, phosphorus solubilizing bacteria and Mycorrhizae are capable of improving the mineral nutrition of plants and enhance the soil fertility in low fertility soils (Shankarappa *et al.*, 2017). Microorganisms such as *Azospirillum* fix atmospheric nitrogen to be taken up by plants; *Pseudomonas fluorescens* control the soil borne pathogens, Mycorrhizae form symbiotic association with plant and supply unavailable phosphorus to plants respectively (Gurumurthy *et al.*, 2014). Phosphorus solubilizing bacteria are capable of solubilizing unavailable form of phosphorus into available form and make it available to plants (Veena *et al.*, 2009; Shankarappa *et al.*, 2012; Shankarappa *et al.*, 2017). Therefore, the effect of biofertilizers on the growth and establishment and nutrient uptake of grafted cashew plants was studied in a pot culture experiment.

Materials and Methods

The experiment on influence of biofertilizers on growth, establishment and nutrient

uptake of grafted mango plants, root stock: totapuri (Bangalora) and scion: alphonso was conducted in poly house at Horticulture Research station, Hogalagere, Kolar during the period 2015-16.

The forty days old rootstocks of mango plants (totapuri), of soft wood tissue were transferred to new polythene bags of size 12 cm x 18 cm that contained potting mixture, soil with farm yard manure and sand in the ratio of 2:1:1 (Soil: FYM; Sand). The potting mixture had pH - 6.95, Organic Carbon - 0.86 %, Available N- 228 kg ha⁻¹, Available P- 35 kg ha⁻¹ and available Potassium 310 kg ha⁻¹. Applied with biofertilizers such as *Azospirillum lipoferum*, Phosphorus solubilizing bacteria (PSB- *Pseudomonas striata*), *Pseudomonas fluorescens*, @ 10 g per graft to form single, dual, triple combinations (10g only) plus vesicular arbuscular mycorrhiza (VAM), soil based biofertilizer @10 g, where ever required. A total of fifteen treatments were formed. The moisture content of soil in poly bags was maintained at field capacity. On the next week the wedge grafting was carried out by using only the soft-wood tissues of the stock and scion (Nagabhushanam, 1985). The seedlings were decapitated at the soft-wood apical region to retain two pairs of leaves and wedge grafting was carried out with a 4-5 cm cleft on the rootstock and a small portion of the inner surface was removed to facilitate a perfect union of the wedge shaped scion. The scion (5 cm length) of variety, alphonso was prepared by shaving a portion of the bark and tissue on either side, inserted in to wedge cut of stock and the union was secured by tying with a 30 cm polythene strip. The top of the scion was covered with a polythene cap to protect the apical portion of the scion from desiccation. The grafts were watered regularly once in a day to maintain moisture content of soil at

field capacity. Polythene wrapping at the union was removed after 75 days after grafting to prevent girdling.

A total of fifteen treatments were tested in the experiment. The details of the treatments are; 1. Uninoculated control, 2. *Azospirillum lipoferum*, 3. Phosphorus solubilizing bacteria, 4. *Pseudomonas fluorescens* 5. vesicular arbuscular mycorrhiza (VAM), 6. *Azospirillum* spp + phosphorus solubilizing bacteria, 7. *Azospirillum* spp + *Pseudomonas fluorescens*, 8. *Azospirillum* spp + Vesicular Arbuscular Mycorrhiza, 9. Phosphorus solubilizing bacteria + *Pseudomonas fluorescens*, 10. Phosphorus solubilizing bacteria + VAM, 11. *Pseudomonas fluorescens* + VAM, 12. *Azospirillum* spp + phosphorus solubilizing bacteria + *Pseudomonas fluorescens*, 13. *Azospirillum* spp + phosphorus solubilizing bacteria + VAM, 14. *Azospirillum* spp + *Pseudomonas fluorescens*+ VAM and 15. *Azospirillum* spp + phosphorus solubilizing bacteria + *Pseudomonas fluorescens* + VAM.

All the biofertilizers were procured from the Department of Agricultural Microbiology, University of Horticultural Sciences, Bangalore. The observations on parameters such as length and girth of the grafted mango plants and number of leaves at an interval of 30 days for 150 days was taken and total biomass, nitrogen and phosphorous content at the end of 150 days of grafting was estimated. The nitrogen and phosphorous of the oven dried plant samples were estimated by microkjeldahl and vanado-molybdate phosphoric acid methods respectively (Jackson, 1973) and the N and P uptake were worked out. Ten replications were maintained to test the per cent establishment and three replications for other parameters and were analyzed statistically using complete randomized design.

Results and Discussion

The evaluation of biofertilizers on growth and establishment of mango grafts var. Alphonso under greenhouse condition showed that the biofertilizers, nitrogen fixer, phosphate solubilizer, biocontrol agent and phosphorous mobilize in the form of *Azospirillum* spp., *Pseudomonas striata*, *Pseudomonas fluorescens* and VAM respectively had increased the growth and nutrient uptake. The growth, establishment and nutrient uptake was significantly superior over un-inoculated control.

All the biofertilizers showed positive influence on the height of the mango grafts (Table 1). The inoculation of biofertilizers, consortia of *Azospirillum*, *Pseudomonas fluorescens* and VAM showed maximum height (68.23 cm) at the end of 150 days after grafting and it was found to be on par with other treatments that received other combinations of three or four beneficial microorganisms.

The increase in height was not prominent in any of the treatments in the initial stages, however, better growth was observed after 90 days after grafting. The un-inoculated control had a height of 50.13 cm i.e. only 18 cm growth from 32.67 cm in 150 days. The consortium of biofertilizer containing three or four beneficial microorganisms gave enhanced growth in terms of height (Akshitha *et al.*, 2014; Shankarappa *et al.*, 2017).

Similarly, the girth of grafted mango showed positive response to biofertilizer inoculation as presented in Table 2. The consortia containing *Azospirillum*, *Pseudomonas fluorescens* and VAM produced maximum girth of 4.07 cm as against the control 3.17 cm observed at the end of 150 days after grafting.

Table.1 Plant height of mango grafts as influenced by biofertilizers

Treatments	Plant height (cm) at different days					
	0	30	60	90	120	150
Uninoculated control	32.67	38.67	40.33	46.67	47.33	50.13
<i>Azospirillum</i> spp. (AZO)	32.67	47.33	48.33	49.00	50.00	58.20
Phosphorus solubilizing bacteria (PSB)	28.33	42.33	46.33	49.33	56.00	59.60
<i>Pseudomonas flourescens</i> (PF)	31.00	41.67	44.00	49.67	52.00	57.47
Vesicular Arbuscular Mycorrhiza (VAM)						
	27.00	34.33	49.67	52.33	54.00	56.27
AZO + PSB	29.00	42.67	48.33	54.00	60.00	66.33
AZO + PF	28.33	40.67	41.33	55.00	57.00	58.50
AZO + VAM	31.67	44.00	49.67	51.00	57.33	59.73
PSB + PF	28.00	34.67	43.67	41.67	48.67	54.00
PSB + VAM	32.00	40.67	45.00	47.33	57.33	63.57
PF + VAM	34.33	41.33	44.33	54.33	55.67	60.10
AZO + PSB + PF	27.67	37.67	49.33	51.00	62.00	63.13
AZO + PSB + VAM	34.33	50.67	51.33	52.67	54.00	57.23
AZO + PF + VAM	34.00	45.00	47.33	56.33	61.00	68.23
AZO + PSB + PF + VAM	26.67	42.00	45.00	57.00	58.67	65.27
S Em ±	5.50	7.84	6.45	7.29	9.05	4.85
CD @ 5%	10.78	15.37	12.65	14.30	17.74	9.50

Table.2 Stem girth of mango grafts as influenced by biofertilizers

Treatments	Plant girth (cm) at different days					
	0	30	60	90	120	150
Uninoculated control	1.87	2.03	2.03	2.53	2.67	3.17
<i>Azospirillum</i> spp. (AZO)	1.60	2.13	2.17	2.37	2.77	3.40
Phosphorus solubilizing bacteria (PSB)	1.73	2.13	2.23	2.77	2.97	3.17
<i>Pseudomonas flourescens</i> (PF)	1.73	2.10	2.13	2.47	2.77	3.20
Vesicular Arbuscular Mycorrhiza (VAM)						
	1.90	2.30	2.57	2.63	2.69	2.97
AZO + PSB	1.83	2.37	2.60	2.63	2.83	3.37
AZO + PF	1.57	1.93	2.03	2.50	2.70	3.03
AZO + VAM	1.73	2.10	2.13	2.47	2.83	2.90
PSB + PF	2.03	2.47	2.67	2.73	2.80	2.96
PSB + VAM	2.00	2.23	2.47	2.73	2.77	3.10
PF + VAM	2.03	2.20	2.50	2.63	2.87	3.20
AZO + PSB + PF	1.80	2.00	2.10	2.67	2.80	3.07
AZO + PSB + VAM	1.97	2.07	2.53	2.83	3.43	3.83
AZO + PF + VAM	2.07	2.23	2.40	2.83	3.20	3.57
AZO + PSB + PF + VAM	2.03	2.20	2.60	2.83	3.63	4.07
S Em ±	0.29	0.31	0.38	0.42	0.38	0.30
CD @ 5%	0.56	0.61	0.75	0.82	0.75	0.59

Table.3 Number of leaves of mango grafts as influenced by biofertilizers

Treatments	No of leaves at different days				
	30	60	90	120	150
Uninoculated control	3.33	5.67	8.33	12.33	13.67
<i>Azospirillum</i> spp. (AZO)	3.00	5.67	9.00	10.33	14.00
Phosphorus solubilizing bacteria (PSB)	3.67	7.00	6.67	8.33	12.00
<i>Pseudomonas flourescens</i> (PF)	4.67	3.33	6.33	10.00	13.33
Vesicular Arbuscular Mycorrhiza (VAM)	4.00	4.33	8.00	9.33	11.00
AZO + PSB	4.67	7.00	8.33	9.00	12.00
AZO + PF	4.67	5.33	9.33	10.33	12.33
AZO + VAM	4.67	6.00	9.33	11.00	13.33
PSB + PF	3.67	6.00	9.00	13.67	16.00
PSB + VAM	3.67	4.00	7.00	14.00	16.33
PF + VAM	5.33	6.67	10.00	14.00	17.33
AZO + PSB + PF	3.00	5.00	6.67	13.33	18.33
AZO + PSB + VAM	3.67	4.00	7.00	11.67	18.67
AZO + PF+ VAM	3.33	6.67	9.67	10.67	17.00
AZO + PSB + PF + VAM	4.67	7.00	7.33	12.33	19.67
S Em ±	0.53	0.62	0.97	1.05	0.69
CD @ 5%	1.03	1.22	1.91	2.07	1.36

Table.4 Total dry weight, N content and P uptake of mango grafts at 150 days

Treatments	Establishment (%)	Total dry wt. (g ⁻¹ plant)	N uptake (mg ⁻¹ plant)	P uptake (mg ⁻¹ Plant)
Uninoculated control	80	12.72	511.00	69.67
<i>Azospirillum</i> spp. (AZO)	100	15.54	573.00	67.67
Phosphorus solubilizing bacteria (PSB)	100	15.37	526.00	76.33
<i>Pseudomonas flourescens</i> (PF)	100	15.43	545.67	72.67
Vesicular Arbuscular Mycorrhiza (VAM)	100	15.83	583.00	75.33
AZO + PSB	100	16.14	577.67	89.00
AZO + PF	100	15.12	595.33	84.33
AZO + VAM	100	14.19	528.00	95.00
PSB + PF	100	14.12	540.67	95.33
PSB + VAM	100	14.84	560.00	97.67
PF + VAM	100	14.83	609.33	97.00
AZO + PSB + PF	100	15.18	614.67	96.00
AZO + PSB + VAM	100	16.25	615.67	99.33
AZO + PF+ VAM	100	16.45	632.00	102.00
AZO + PSB + PF + VAM	100	16.50	637.33	107.33
S Em ±		0.08	6.18	2.40
CD @ 5%		0.15	12.18	4.69

The biofertilizers consortia of three or four organisms were significantly superior over consortia of two organisms or biofertilizer of single organisms (Veena *et al.*, 2009; Shankarappa *et al.*, 2017). Although, a steady increase in girth was observed in all the treatments, it was more at the later part of the growth, after 90 days of grafting and the growth was superior in biofertilizers treated grafts.

The effect of biofertilizer inoculation on number of leaves produced in grafted mango was significantly superior in the consortia of three or four organisms (Table 3). The treatment, *Azospirillum*, PSB, *Pseudomonas fluorescens* and VAM (19.67) showed maximum number of leaves which is superior over other treatments of combinations of *Azospirillum*, *Pseudomonas fluorescens* and VAM inoculation. The control treatment produced only 13.67 numbers of leaves at the end of 150 days after grafting. Here again the number of leaves had increased after 90 days of grafting and all the biofertilizers could increase the number of leaves.

Establishment of grafted plants, biomass output and nutrient uptake was observed to be high in biofertilizers treated mango grafts (Table 4). The success of grafted plants to establish was 100 percent in biofertilizer treated grafts as against 80 percent observed in control. All the biofertilizers irrespective of their application as single, dual, triple or four organisms in a consortium produced better establishment. The establishment of grafts, increased graft height, girth and number of leaves due to biofertilizer inoculation had reflected on biomass output (Shankarappa *et al.*, 2012; Vijendrakumar *et al.*, 2014; Shankarappa *et al.*, 2017). The total dry matter content was high in biofertilizer treated grafts. Highest biomass of 16.50 g was observed in the treatment

Azospirillum, PSB, *Pseudomonas fluorescens* and VAM followed by was *Azospirillum*, *Pseudomonas fluorescens* and VAM inoculation (16.45). These treatments were on par with other treatments of triple organisms inoculations. Similarly, the nitrogen uptake (637.33 mg) and phosphorous uptake (107.33 mg) per plant was found to be highest in the plants treated with consortia of biofertilizers that contained all the four organisms and it was on par with consortia that contained combinations of three organisms. The enhanced height, girth and number of leaves of mango grafts and better nutrient uptake and biomass growth, observed due to inoculation of single or multiple biofertilizers is attributed to the production of various growths promoting hormones by the beneficial microorganisms such as IAA, Gibberellic acid, and cytokinins (Veena *et al.*, 2009; Shankarappa *et al.*, 2012) as well as the availability of fixed nitrogen by *Azospirillum*, phosphate solubilization by *Pseudomonas striata* (Akshitha *et al.*, 2014; Vijendrakumar *et al.*, 2014), phosphate mobilization by the mycorrhizal fungi (Gurumurthy *et al.*, 2014), biocontrol activity of *Pseudomonas fluorescens* (Akshitha *et al.*, 2014; Vijendrakumar *et al.*, 2014; Shankarappa *et al.*, 2017) and the synergistic growth promotion by the consortia of biofertilizers used.

The experiment on effect of biofertilizers establishment and nutrient uptake of mango grafted plants showed that the inoculation of consortium of three biofertilizers; *Azospirillum*, *Pseudomonas fluorescens* and VAM was efficient in getting maximum plant height (68.23 cm), stem girth (3.83 cm) and number of leaves (18.67 no.). Also the inoculation of consortia of these three biofertilizers showed better nutrient uptake and growth at the end of 150 days in nursery. The inoculation of consortium of

three biofertilizers was found to be on par with inoculation of consortium containing four biofertilizers but found to be superior over dual inoculation, single inoculation and over control.

References

- Akshitha, J., Umesha, K. and Shankarappa, T.H. 2014. Effects of type of cutting, IBA and bioinoculants on rooting in madhunashini (*Gymnema sylvestre* Retz.). *J. Hort. Sci.*, 9(1):94-97.
- Gurumurthy, S. B., Patil S. V., Vidyavathi, K. B., Lokesh, M. S., Shankrappa, T.H. and Bellakki. M.A. 2014, Performance of local isolates of arbuscular mycorrhizal (AM) fungi on growth and yield of chilli (*Capsicum annuum* L.) grown in black clayey soil. *Int.J.Curr.Microbiol.App.Sci.*, 3(9) 404-408.
- Jackson, M. L. 1967. Seed chemical analysis, Prentice Hall of India PVT. Ltd., New Delhi, pp183-192.
- Mehta, I. 2017. History of Mango - King of Fruits, *Int. J Engg. Sci. Invention* 6(7): 20-24.
- Nagabhushanam, S. 1985. Vegetative propagation in cashew - Review of work done at vitla. *Acta Hort.*108:59-65.
- National mango database, 2015. mangifera.res.in/indianstatus.php.
- Shankarappa, T.H., Gurumurthy, S.B., Patil S.V. and Lokesh, M.S. 2012. Influence of phosphorus enriched biogas spent slurry (BSS) on growth and yield of sunflower (*Helianthus annuus*). *Int. J. Pl Sci.*, 7 (2):253-258.
- Shankarappa, T.H., S. K. Mushrif, B. Subramanyam, A. Sreenatha, B. N. Maruthi Prasad and N. Aswathanarayana Reddy., 2017. Effect of biofertilizers on growth and establishment of cashew grafts under nursery condition. *Int. J. Curr. Microbiol. App. Sci.* 6(8): 1959-1965.
- Veena, S.C., Alagawadi, A.R., Shankarappa, T.H. and Krishnaraj, P.U. 2009. Development of inoculums consortia for improved performance: II. Impact on growth and nutrient uptake of Sorghum (*Sorghum bicolor* L. Moench). *J. Soil Bio Ecol.* 29 (1&2): 52-59.
- Vijendrakumar, R. C., Sreeramu, B. S., Shankarappa, T.H., Santhosh, K.V., Mallikarjuna Gowda, A.P. and Umesha, K. 2014. Effect of liquid bio fertilizers on growth, yield and survival of seedlings in garden rue (*Ruta graveolens* Linn.). *Plant Archives*, 14(1): 171-175.