



## Original Research Article

### Isolation of wood rot fungi from Maruthamalai hills Western Ghats area of south India

M.Meenakshisundaram<sup>1\*</sup> and C.Bharathiraja<sup>2</sup>

<sup>1</sup>Post Graduate & Research, Department of Biotechnology, Nehru Memorial College, (Autonomous), Puthanampatti, Trichy-621 007, Tamilnadu, India

<sup>2</sup>Department of Environmental Biotechnology, Bharathidasan University, Trichy, Tamilnadu, India

\*Corresponding author e-mail: [meenakram06@yahoo.co.in](mailto:meenakram06@yahoo.co.in)

#### ABSTRACT

##### Keywords

Basidiomycete;  
Isolation;  
Western Ghat;  
wood rot  
fungi.

An attempt was made to isolate the wood rot fungi from the different host plants at maruthamalai hill forest area. This is a located in western ghatsof southern India. There are 30 wood rot fungal samples were observed . This is first report of kind from maruthamalai hill forest among the white rot fungi. These fungi might have played very important role in sustaining in forest ecosystem in maruthamalai hill forest of western ghats, South India. From the above isolated thirty wood rot fungi samples were indentified and screening the best wood rot fungi for further studies.

#### Introduction

Wood is formed of three main constituents, cellulose, hemicelluloses and lignin. Lignins constitute the second most abundant group of biopolymers in the biosphere (Karthikeyan, *et al.*, 2005) Lignin is highly resistant towards chemical and biological degradation, and confers mechanical resistance to wood (Martinez, *et al.*, 2005). The plant material contains 25-30% of lignin, which gives plants their structural integrity and provides protection from pests and pathogens (Orth, *et al.*, 1993). Lignins are polymers consisting of phenylpropane units in which the aromatic rings are substituted with guaiacyl propane units, syringylpropane units or parahydroxyphenylpropane units (Ralph *et al.*, 2004). Wood rotting fungi are an

important component of forest ecosystems (Wang, *et al.*, 2011). White rot fungi belong to the order basidiomycetes that participates in the biodegradation of lignin in nature, which is essential for global carbon recycling (Siripong, *et al.*, 2009). White rot fungi can degrade lignin and a range of environmental pollutants by many of their extra cellular ligninolytic enzymes (Selvam, *et al.*, 2011).

White rot fungi possess a number of advantages that can be exploited in bioremediation systems, since the key components of their lignin-degrading system are extracellular, these fungi can degrade an extremely diverse range of very persistent or toxic environmental

pollutants (Silvia, *et al.*, 2008). The ability of fungi to degrade lignocellulosic materials is due to their highly efficient enzymatic system (Sanchez, 2009). Microbial degradation of lignin has of cellulose are extremely attractive for use in biological pulping processes, to improve the digestibility of highly lignified received considerable attention in recent years. Fungi that selectively remove lignin without loss of appreciable amounts plant industrial products (Kirk, *et al.*, 1980). White rot fungi emerged as a promising group for biotechnological applications, especially in bioremediation (Reddy, *et al.*, 2011). In the present study isolation of 30 samples of wood rot fungi from different host plants have been carried out Maruthamalai hill

## **Materials and Methods**

### **Collection and Isolation**

Thirty samples of fruiting bodies of the wood rot fungi were collected from decayed wood and living trees of ofmaruthamalai hill, western ghats, Tamilnadu. The samples were marked with information such as collection number with names, procurement location and date of collection. Fruit bodies were wrapped in paper bags and brought to the laboratory. The collection sight is situated in the 76° 55E and 110.5 ° N.

It receives rainfall of about 450mm per year with humidity and even temperature. The soil condition was generally shallow with sandy loam texture, rocky substratum was available at steepy areas (Paulsamy, 2012).

The wood rot fungal sporocarps were photographed at the sight using digital camera. The various wood rot fungi and

their host were noted with the help of local tribal people. The fungal sporocarps were carefully in two sets of samples (Mueller, *et al.*, 2004).

One set of the sample collected were preserved for culturing the mycelia for further studies. The second set of the samples were used for the study of morphological (Colour, shape, and odour) characters, microscopical characters and the biochemical identification according to the method of walting, (1971).The fungal growth was cut sterilized with 1% mercuric chloride solution, repeatedly washed with sterile distilled water (Walting, 1971) and inoculated on 2 % malt agar medium in petriplates. The fungal growth which occurred on the plates were subcultured and maintained in malt agar slants.

## **Results and Discussion**

### **Collection of the Samples**

In the present study Thirty samples were collected from various location of maruthamalai hill area of Tamilnadu in South India. General macrocharacters of the fruit body including colour of different tissues and type of rot were noted in the field. The isolation of wood rot fungi and their Character are presented in Table-1

## **Acknowledgement**

This research work was funded by Science Engineering Research Board, (SERB), New Delhi, Government of India FAST Track scheme for research project, No.SR/FT/LS-38/2012. The corresponding author thank (SERB) for their constant financial support and guidance rendered throughout the period of study.

**Table.1** Detailed of wood rot fungi collected from maruthamalai

S. No	Strain No	Mushroom vernacular name	Mushroom character	Host botanical name
1.	M1	Marakaalaan	Non edible	<i>Acacia lucophloea</i>
2.	M2	Marakaalaan	Edible	<i>Acacia nilotica</i>
3.	M3	Vellaikaalaan	Edible	<i>Agave americanum</i>
4.	M4	Marakaalaan	Edible	<i>Albizziaamericanum</i>
5.	M5	Unknown	Non edible	<i>Albizzialebeck</i>
6.	M6	Unknown	Non edible	<i>Atlantiamonophylla</i>
7.	M7	Unknown	Non edible	<i>Azadirachtaindica</i>
8.	M8	Unknown	Non edible	<i>Bambusaarundinacea</i>
9.	M9	Marakaalaan	Edible	<i>Bouhiniavariegata</i>
10.	M10	Marakaalaan	Non edible	<i>Cassia fistula</i>
11.	M11	Marakaalaan	Non edible	<i>Cassia siamea</i>
12.	M12	Marakaalaan	Non edible	<i>Delonixregia</i>
13.	M13	Marakaalaan	Non edible	<i>Ficusbengalensis</i>
14.	M14	Unknown	Non edible	<i>Pongamiapinnata</i>
15.	M15	Kuyaankaalaan	Edible	<i>Santalum album</i>
16.	M16	Marakaalaan	Non edible	<i>Tamarindusindica</i>
17.	M17	Marakaalaan	Non edible	<i>Tectonagrands</i>
18.	M18	Marakaalaan	Non edible	<i>Terminaliarjuna</i>
19.	M19	Marakaalaan	Non edible	<i>Thevitaperviana</i>
20.	M20	Marakaalaan	Non edible	<i>PolyalthialongifoliaS.nn</i>
21.	M21	Marakaalaan	Non edible	<i>MurrayaKoenigii</i>
22.	M22	Seenikallan	Non edible	<i>Azadirachtaindica</i>
23.	M23	Marakaalaan	Non edible	<i>Mangiferaindica</i>
24.	M24	Marakaalaan	Non edible	<i>Moringapterygosperma</i>
25.	M25	Yaanaimithi	Soil - Edible	<i>Dalbergiasissoo</i>
26.	M26	Sarugumutti	Soil - Edible	<i>Sesbaniagrandiflora</i>
27.	M27	Neeikaalaan	Soil - Edible	<i>Cassia roxburghii</i>
28.	M28	Yaanaikaathu	Soil - Edible	<i>Acacia auriculiformis</i>
29.	M29	Marakaalaan	Edible	<i>Ficusbenghalensis</i>
30.	M30	Vellaikaalaan	Edible	<i>Cocosnucifera</i>

## References

- Karthikeyan, S., K. Ramasamy and Prabhakaran, J. 2005. Conversion of industrial solid wastes to value added products by Filamentous Bacteria. *Madras.Agric.J.* 92: 515-522.
- Kirk, T.K., T. Higuchi and Chang, R. 1980. Lignin biodegradation, summary and perspectives. *Lignin biodegradation, microbiology., chemistry and potential applications*, CRC Press. Inc. Boca Raton., Fla.1: 235-244.
- Matrinez, T.A., M.S. Francisco, R.J. Duenas, P. Ferrerira, S. Camarero, F. Guillen, M.J. Matrinez, A. Gutierrez and Rio, J.C. 2005. Biodegradation of lignocellulosics: microbial., chemical and enzymatic aspects of fungal attack of lignin, *Res Rev.* 8: 195-204.
- Mueller, G., G.F. Bills and Foster, M.S. 2004. Biodiversity of fungi: Inventory and monitoring methods. Elsevier Academic press. 1: 23-26.
- Orth, B.A., D.J. Royle and Tien, M. 1993. Ubiquity of Lignin-Degrading Peroxidases among Various Wood-Degrading Fungi. *Appl. Environ. Microbiol.* 3: 4017-4023.
- Paulsamy, S., 2012. Maruthamalai hills of western Ghats, Coimbatore District, Tamil Nadu. A Potential ecosystem for medicinal plants. *J. Res.Plant sci.* 1: 12-26.
- Ralph, J., K. Lundquist, G. Brunow, F. Lu, H. Kim, P.F. Schatz, J.M. Martia, R.D. Hatfield, S.A. Ralph, J.H. Christensen and Boerjan W. 2004. Lignins, Natural polymers from oxidative coupling of 4-hydroxyphenylpropanoid. *Phytochem. Rev.* 3: 29-60.
- Reddy, C.A., and Mathew, Z. 2001. Bioremediation potential of white rot fungi. *Fungi in bioremediation*, Gadd GM. Cambridge.,UK. Cambridge University Press. pp. 1-472.
- Sánchez, C., 2009. Lignocellulosic residues, Biodegradation and bioconversion by fungi. *Biotechnol. Adv.* 27: 185-194.
- Selvam, K., M. ShanmugaPriya and Arungandhi, K. 2011. Pretreatment of Wood Chips and Pulps with *Thelephora Sp.* to reduce Chemical Consumption in Paper Industries, *Inter. J. Chem.Tech. Res.* 3: 471-476
- Silvia, F., and Naresh, M. 2008. Impact of *Trametes versicolor* and *Phanerochaete chrysosporium* on differential breakdown of pesticide mixtures in soil microcosms at two water potentials and associated respiration and enzyme activity, *Intbiodegrad. biodegrad.* 62:376-383.
- Siripong, P., B. Oraphin, T. Sanro and Duanporn, P. 2009. Screening of Fungi from Natural Sources in Thailand for Degradation of Polychlorinated Hydrocarbons. *Am. Eurasian. J. Agric. Environ.Sci.* 5: 466-472.
- Walting, C., 1971. Chemical tests of agaricology. In: *methods of microbiology*, ed, Academic press, New York. pp.567- 597.
- Wang, B., K.B. Cuil, H.J. Li, P. Du and Jia, B.S. 2011. Wood rotting fungi from eastern China. 5. polypore diversity in Jiangxi province. *Ann. Bot. Fennici.* 48: 237-246.
- Watling, R., 1971. Basidiomycetes, Homobasidiomycetidae. In, *Methods in Microbiology.*, (ed. Booth., C.). Academic press., London and New York. pp. 219.