



Original Research Article

In Vitro Study of Lysis of Cell Wall Preparation from *Phomopsis vexans* by Lytic Enzyme from Some Biocontrol Agents

S. K. Ghosh*, S. Pal, S. Banerjee and N. Chakraborty

Molecular Mycopathology Lab. Dept. of Botany, Ramakrishna Mission Vivekananda Centenary College, Rahara, Kolkata -700118, West Bengal, India

*Corresponding author

ABSTRACT

Keywords

Lytic enzyme,
Biocontrol
agents,
P. vexans

Cell wall was prepared from *P. vexans* (Sacc.&Syd.), dangerous *Phomopsis* blight and fruit rot disease of brinjal (*Solanum melongenum* L). *In vitro* the potentiality of lytic enzyme production by three biocontrol agents (*Trichoderma viride* -1, *T. harzianum* -1 and *Beauveria bassiana*) was tested in mineral medium. In our experiment all three antagonists or biocontrol agents showed their ability to secrete lytic enzyme. Out of them the enzyme of *T. harzianum*-1 lysed 67% of the cell wall after 48 hrs, it was followed by *T. viride* I (55%) and *Beauveria bassiana* (47%). Moreover, within 24 hour, at 25°C, the lysis of enzyme preparation of *T. harzianum* -1 is maximum (35%) followed by 20°C (29%), 30°C (20%) and 15°C (20%) and it indicated that at 25°C, the efficacy of enzyme secreted by *T. harzianum* -1 is optimum. It revealed that biocontrol agents during interaction with this pathogen may have secreted this enzyme to suppress or kill the pathogen.

Introduction

Lytic enzymes among the diverse metabolites secreted by microorganisms can interfere with pathogens' growth and/or activities and these enzymes are important weapons for killing pathogenic fungi or pathogens. Lytic enzymes can hydrolyze a wide variety of polymeric compounds, including chitin, proteins, cellulose, hemicellulose, and DNA. Expression and secretion of these enzymes by different biocontrol agents cause in the suppression of plant pathogen activities directly. Control of *Sclerotium rolfsii* by *Serratia marcescens* appeared to be mediated by chitinase

expression (Ordentlich *et al.*, 1988) and a β -1,3-glucanase contributes significantly to biocontrol activities of *Lysobacter enzymogenes* strain C3 (Palumbo *et al.*, 2005). While they may stress and/or lyse cell walls of living organisms, these enzymes generally act to decompose plant residues and nonliving organic matter. Currently, it is unclear how much of the lytic enzyme activity that can be detected in the natural environment represents specific responses to microbe-microbe interactions. It seems more likely that such activities are largely indicative of the need to degrade

complex polymers in order to obtain carbon nutrition. Furthermore, some products of lytic enzyme activity may contribute to indirect disease suppression. For example, oligosaccharides derived from fungal cell walls are known to be potent inducers of plant host defences. Interestingly, *Lysobacter enzymogenes* strain C3 has been shown to induce plant host resistance to disease (Kilic-Ekici and Yuen, 2003), though the precise activities leading to this induction are not entirely clear.

The quantitative contribution of any and all of the above compounds to disease suppression is likely to be dependent on the composition and carbon to nitrogen ratio of the soil organic matter that serves as a food source for microbial populations in the soil and rhizosphere. However, such activities can be manipulated so as to result in greater disease suppression.

For example, in post-harvest disease control, addition of chitosan can stimulate microbial degradation of pathogens similar to that of an applied hyperparasite (Benhamou, 2004). Chitosan is a non-toxic and biodegradable polymer of beta-1,4-glucosamine produced from chitin by alkaline deacylation.

Amendment of plant growth substratum with chitosan suppressed the root rot caused by *Fusarium oxysporum* f. sp. *radicis-lycopersici* in tomato (Lafontaine and Benhamou, 1996). Although the exact mechanism of action of chitosan is not fully understood, it has been observed that treatment with chitosan increased resistance to pathogens. Therefore, the main objectives of this work are to prepare cell wall from some plant pathogens, to produce lytic enzyme from biocontrol agent and finally to evaluate of enzyme action or cell wall and enzyme interaction.

Material and Methods

Preparation of cell wall of the fungal pathogen- *Phomopsis vexans*

The pathogen growing on PDA slants was harvested after 20 days of growth in 5 ml of distilled sterile water from each slant. An aliquot of 25 ml of pathogen suspension was obtained from 5 slants. They were crushed and homogenized in a homogenizer (A. HT Philadelphia, U.S.A). It was washed by repeated centrifugation (3000 rpm) with 0.1M NaCl, 0.1M acetate buffer (pH= 5.5) and distilled water until the cells were free from cytoplasmic materials. The wall was inactivated by heating at 100°C for 30 minutes in a boiling water bath. An aliquot of 0.2µM Na₂NO₃ was then added to keep the preparation sterile.

Preparation of enzyme from biocontrol agents

Trichoderma viride-1, *Trichoderma harzianum*-1, & *Beauveria bassiana* were grown separately in 250 ml conical flasks containing 100 ml mineral medium for 8 days at 25 ±1°C.

Mineral medium

Glucose (anhydrous):10 g; Ammonium tartarate:2g; K₂HP0₄:1 g; MgSO₄·7H₂O:0.5g; Trace element solution:1 ml;pH:5.5.

The trace element solution in (mg/L) : Na₂B₄O₇, 10H₂O, 100; ZnSO₄, 7 H₂O, 70; FeSO₄, 7H₂O, 50; CuSO₄, 5H₂O, 10; MnSO₄, 4H₂O, 10; (NH₄)₆ Mn₇O₂₄, 4H₂O, 10.

The culture filtrate was separated from the mycelium by filtration through Whatman Filter No. 3. The filtrate was centrifuged at 3000 rpm for 30 minutes. The culture filtrate

was then dialysed twice against two changes of double distilled water for 48 hrs at 4°C. This resultant preparation was taken as that of enzyme.

Measurement of lysis

One mg. of cell wall preparation was taken in 1 ml. of 0.05 M borate- citrate-phosphate buffer (pH 5.5). This was inoculated for different times (12 hr, 24hr 36 hr, 48 hr) and temperature (5°C, 10°C, 15°C, 20°C, 25°C, 30°C and 35°C) with 1 ml enzyme preparation. Turbidity was measured in Nephelometric turbidity units (NTU) by Nephelometric method with the help of a Nephelometer (Systronic, India). The sample was diluted with one or more volumes of turbidity free water until turbidity level fell down within limits of 30 to 40 NTU. Turbidity in original sample was calculated from the turbidity of diluted sample and the dilution factor was known. The sample was then transferred to the turbidometer tube and direct reading was taken on the scale. The calculation was done by using this formula:

Nephelometric Turbidity Units = $A \times (B+C)/C$ where A= NTU found in diluted sample, B= Volume of dilution water in ml; C = sample Volume taken for dilution in ml.

The percentage loss in turbidity indicates the measure of lysis of the cell wall of pathogens by enzymes of different antagonists or biocontrol agents.

The boiled enzyme (30 minutes) was taken as control for the experiment.

Result and Discussion

The data presented in table 1 revealed the degree of cell wall lysis of *P.vaxans* by enzymes of biocontrol agents (*T. harzianum* -1, *T. viride* -1 and *B. bassiana*). The enzyme of *T. harzianum* -1 lysed 68% of the cell wall after 48 hrs, it was followed by *T. viride* -1 (55%) and *Beauveria bassiana* (47%). The involvement of enzyme in mechanism of mycoparasitism of mycoparasites is recorded by many workers (Elad *et al.*, 1983). Panchenari and Dix (1980) observed that *Gliocladium roseum* parasitized on *B. alli* by glucanase enzyme which degrades glucans of the cell wall. Elad *et al.* (1983) revealed lysed sites and penetration holes at the area of contact of *R. solani* hyphae parasitized by *Trichoderma* spp. They detected with scanning electron microscope and fluorescence microscope high amount of secretion of β -1, 3 glucanase and chitinase in dual agar cultures where *T. harzianum* parasitized *S. rolfsii*.

Table.1 Degree of lysis (%) of the cell wall preparation of the pathogen by different biocontrol agents at 25±10°C

Antagonists	*Percentage of lysis of cell wall <i>P. vexans</i>			
	12 hr	24hr	36 hr	48hr
<i>Trichoderma viride</i> -1	17	25	37	55
<i>T. harzianum</i> -1	25	36	55	68
<i>Beauveria bassiana</i>	15	24	35	47

*percentage mean data of three replicas

Table.2 Effect of different temperatures on the lyses cell wall by enzyme preparation of *T. harzianum*-1

Incubation temperature °C	Lysis (%) of <i>P. vexans</i> cell wall preparation measured after 24 hour of incubation
5	00
10	05
15	20
20	29
25	35
30	20
35	16

The enzyme of *T. harzianum* rapidly digested (86%) the cell wall of *R. solani* within 25 hrs (Srivastava and Singh, 2000). Lytic enzyme of *T. harzianum* digested 76% of the cell wall preparation of *A. rabiei* (Chakraborty *et al.*, 2008). Ghosh *et al.* (2015) recorded *T. harzianum* -1 secreted lytic enzyme and lysed 45% of cell wall preparation of *P. digitatum*. In this experiment, enzymes of *T. harzianum* 1 showed the same trend. The table 2 showed that within 24 hour, at 25°C, the lysis of enzyme preparation of *T. harzianum*-I is maximum (35%) followed by 20°C (29%) 30°C (20%) and 15°C (20%) and it indicated that at 25°C, the efficacy of enzyme secreted by *T. harzianum*-I is optimum.

In conclusion, all three biocontrol agents (*T. harzianum*-1, *T. viride* -1 and *B. bassiana*) produced lytic enzymes and lysed cell wall preparation of *P. vexans*. It gives clue that biocontrol agents during interaction between this pathogen may have secreted this enzyme to suppress or kill the pathogen.

Acknowledgement

Authors are grateful to the Principal of RKMVC College for all kind of lab facilities.

Reference

- Benhamou, N. 2004. Potential of the mycoparasite, *Verticillium lecanii*, to protect citrus fruit against *Penicillium digitatum*, the causal agent of green mold: A comparison with the effect of chitosan. *Phytopathology*, 94: 693–705.
- Chakraborty, N., Ghosh, S.K., Sengupta, C. 2008. Study of lysis of cell wall preparation of *Ascochyta rabiei* (pass) Labr. by different antagonists. *Indian J. Bot. Res.*, 4(3&4): 465–468.
- Elad, Y., Chet, I., Henis, Y. 1983. Parasitism of *Trichoderma* spp. on *Rhizoctonia solani* and *Sclerotium rolfsii* scanning microscopy and fluorescence microscopy. *Phytopathology*, 73: 85–88.
- Ghosh, S.K., Pal, S., Banerjee, S. 2015. Study of cell wall preparation of *Penicillium digitatum* and lytic enzyme from some biocontrol agents and their interaction. *Int. J. Curr. Res.*, 7(8): 19383–85.
- Kilic-Ekici, O., Yuen, G.Y. 2003. Induced resistance as a mechanism of biological control by *Lysobacter enzymogenes* strain C3. *Phytopathology*, 93: 1103–1110.

- Lafontaine, P.J., Benhamou, N. 1996. Chitosan treatment: an emerging strategy for enhancing resistance of greenhouse tomato plants to infection by *F. oxysporum* f.sp *radicis-lycopersici*. *Biocontrol Sci. Technol.*, 6: 111–124.
- Ordentlich, A., Elad, Y., Chet, L. 1988. The role of chitinase of *Serratia marcescens* in biocontrol of *Sclerotium rolfsii*. *Phytopathology*, 78: 84–88.
- Palumbo, J.D., Yuen, G.Y., Jochum, C.C., Tatum, K., Kobayashi, D.Y. 2005. Mutagenesis of beta-1,3-glucanase genes in *Lysobacter enzymogenes* strain C3 results in reduced biological control activity toward *Bipolaris* leaf spot of tall fescue and *Pythium* damping-off of sugar beet. *Phytopathology*, 95: 701–707.
- Panchenari, A., Dix, N.J. 1980. Production of toxins and wall degrading enzymes by *Gliocladium roseum*. *Trans. Brit. Mycol. Soc.*, 74: 561–566.
- Srivastava, A.K., Singh, K.R.K. 2000. Extent of lysis of *Rhizoctonia solani* Kuhn. cell wall preparation by different hyperparasites. *J. Mycopathol.*, 38(2): 129–131.